

Tamarindus indica seed extracts inhibit mouse microsomal 5 α -reductase *in vitro*

Gerwin Louis Tapan DELA TORRE^{1*} , Richelle Ann Mallapre MANALO^{1,2} , Erna Custodio AROLLADO^{1,2} 

¹ Institute of Pharmaceutical Sciences, National Institutes of Health, University of the Philippines Manila, Manila, Philippines

² Department of Pharmacy, College of Pharmacy, University of the Philippines Manila, Manila, Philippines

* Corresponding Author. E-mail: gtdelatorre@up.edu.ph (G.L.T.D.); Tel. +63-2-7577-2001.

Received: 02 March 2023 / Revised: 15 April 2023 / Accepted: 18 April 2023

ABSTRACT: *Tamarindus indica* seeds have important roles in the field of health sciences. The seeds are used in traditional medicine due to their beneficial effects on urologic disorders, including benign prostatic hyperplasia. The present study evaluates the inhibitory activity of *T. indica* seed extracts against 5 α -reductase (5 α R). This enzyme is responsible for the conversion of testosterone to dihydrotestosterone, which stimulates prostatic enlargement. Seeds were separately extracted with ethanol, ethyl acetate, and hexane. The inhibitory activity of the extracts was assessed in the presence of testosterone and 5 α R from mouse liver. Unchanged testosterone was quantified by high-performance liquid chromatography with photodiode array detector. Inhibition was calculated based on the remaining testosterone and was expressed as dutasteride equivalent 5 α R inhibitory activity (DE; μ g dutasteride equivalent per mg extract). The most active extract underwent phytochemical profiling by ultra-performance liquid chromatography quadrupole time-of-flight mass spectrometric analysis. The results showed that ethyl acetate extract was the most active 5 α R inhibitor with a DE value of 153.31 ± 10.60 . Fifteen phytochemicals from the classes of tocopherol, fatty acid, phytosterol, saponin, terpenoid, and phenolic compounds were identified, which may be partly responsible for the inhibitory activity of the extract. The findings indicate that *T. indica* seeds can be a source of 5 α R inhibitors.

KEYWORDS: 5 α -reductase; benign prostatic hyperplasia; dihydrotestosterone; *Tamarindus indica*; testosterone.

1. INTRODUCTION

Benign prostatic hyperplasia (BPH) is a common condition in middle-aged to elderly men, characterized by an enlargement of the prostate gland as a result of an increase in epithelial and stromal cells [1]. This causes the urethra to be compressed, causing several symptoms in the lower urinary tract [2]. Although the precise molecular etiology of BPH is complex and not fully understood, androgens have been linked to this condition because the growth and development of the prostate are androgen-dependent [3, 4]. In men, testosterone accounts for approximately 90% of the circulating form of androgen [5]. It is irreversibly converted to dihydrotestosterone (DHT) by the action of microsomal enzyme 5 α -reductase (5 α R), which is intrinsically more potent in stimulating the growth of the prostate [6]. The potency may be related to the superior affinity and selectivity of DHT over testosterone for androgen receptors [7]. The accumulation of DHT within the prostate acts as the hormonal mediator of hyperplasia [8]. Studies have concluded that individuals with low DHT levels due to reduced expression of 5 α R have small prostates. Moreover, those with 5 α R deficiency do not have palpable prostates, and the occurrence of BPH has not been reported [9, 10].

With the knowledge that 5 α R plays a pivotal role in the production of DHT, and that DHT is affirmatively linked to the pathologic growth of the prostate, therapeutic inhibitors of the enzyme have been identified [11, 12]. Dutasteride is a 5 α R inhibitor drug that markedly suppresses DHT serum levels by effectively preventing the conversion of testosterone to DHT [13]. However, its use is limited by its multiple side effects such as impotence, erectile dysfunction, and decreased libido. To answer this problem, plant-based alternatives are being studied because they are believed to be more effective and well-tolerated by the body than their synthetic counterparts [14, 15].

Tamarindus indica seeds have been used in healthcare since time immemorial [16]. The seeds are considered to be agri-industrial by-products but were found to be applicable as food additives and pharmaceutical excipients. The plant is also one of the most widely used herbal medicines against urologic

How to cite this article: Dela Torre GLT, Manalo RAM, Arollado EC. *Tamarindus indica* seed extracts inhibit mouse microsomal 5 α -reductase *in vitro*. J Res Pharm. 2023; 27(5): 1989-1997.

problems due to its wide geographic distribution [17]. For instance, the decoction of *T. indica* seeds has been traditionally used in the treatment and management of BPH [18]. With the view of providing scientific support for its use in BPH, we evaluated the *in vitro* 5 α R inhibitory activity of *T. indica* seed extracts and qualitatively analyzed the phytochemicals present in the active extract.

2. RESULTS AND DISCUSSION

Three *T. indica* crude seed extracts, namely, *T. indica* ethanol extract (TIEE), *T. indica* ethyl acetate extract (TIEAE), and *T. indica* hexane extract (TIHE) were obtained via maceration. All extracts were dark green, with sticky consistency and semisolid appearance. The TIEE showed the highest yield of 2.35% followed by TIEAE (0.95%) and TIHE (0.82%). This indicates that polar compounds occur abundantly in the seeds of *T. indica*.

The optimized method for the assay of testosterone is summarized in Table 1. The α,β -unsaturated ketone in the A-ring of testosterone absorbs maximum UV light at 240 nm [19], thus this wavelength was used. Previous studies reported an isocratic elution of the mobile phase composed of methanol and water for the analysis of testosterone. The use of a binary solvent mixture has been well reported to be efficient in the identification of analytes [20] and the appropriate solvent ratio was probed to obtain a chromatogram that presents a better separation of analyte and internal standard (IS). For the assay of testosterone, a suitable IS was selected from a pool of compounds, namely propylparaben, ursodeoxycholic acid, and cholecalciferol. Individually, all three compounds absorb UV at 240 nm, with peaks eluting at around 2 min. However, with testosterone, clear peak separation was only observed using propylparaben. In addition, propylparaben was stable and non-reactive with testosterone, hence it was chosen as the IS for the study. A flow rate of 1.00 ml/min provided early elution of testosterone and propylparaben. In the study, a high injection volume of 10 μ l produced broad peaks, resulting in bad separation. This was improved by lowering the injection volume to 5 μ l. An elevated column temperature at 50°C was also set to improve column efficiency, lessening the effects of peak broadening [21]. Given these conditions, propylparaben and testosterone gave retention times of 2.1729 \pm 0.0037 and 3.1045 \pm 0.0061 min, respectively. A representative chromatogram is presented in Figure 1.

Table 1. Optimized HPLC-PDA method for the detection and quantification of testosterone.

Parameter	Optimized Condition
UV detection	240 nm
Mode of elution	Isocratic
Mobile phase composition	Methanol:Water (90:10)
Internal standard	Propylparaben
Flow rate	1.00 ml/min
Injection volume	5 μ l
Column temperature	50°C




Figure 1. Representative chromatogram of propylparaben (IS) and testosterone.

The use of the mouse liver as a source of 5 α R in conjunction with high-performance liquid chromatography with photodiode array detector (HPLC-PDA) for the detection and quantification of unchanged testosterone was recognized to be useful for the analysis of 5 α R inhibitory activity of various compounds. Prior to the HPLC-PDA analysis, the protein content of the resulting pink microsomal pellets was quantified using the modified Lowry method. It is a combination of reactions from the Biuret method where

copper ions react with peptide bonds along the protein chain and the Folin-Ciocalteu reagent reacting with the ring structure of aromatic amino acids within the protein [22]. These reactions produced a stable, dark blue complex. Based on the bovine serum albumin curve, the microsomal pellet had a protein concentration of 840 $\mu\text{g}/\text{ml}$.

The microsomal pellets isolated were used for the determination of the 5 α R inhibitory activity of the extracts. Dutasteride gave a half-maximal inhibitory activity (IC_{50}) of $109.75 \pm 4.53 \mu\text{g}/\text{ml}$. The inhibitory equation (Eq. 1) of dutasteride was:

$$\text{(Eq. 1) } y = 0.1856x + 22.622 \text{ (} R^2 = 0.9975 \text{)}$$

with y representing the % inhibition and x representing the concentration of dutasteride. This equation was used to calculate the 5 α R inhibitory activities of the extracts expressed as dutasteride equivalent 5 α R inhibitory activity (DE; μg dutasteride equivalent per mg extract). The DE values of extracts are summarized in Table 2. A high DE value signifies a strong 5 α R. The varied inhibitory potential of the extracts may be attributed to the amount and classes of phytochemicals present. Based on the study, the highest activity shown by TIEAE suggests that moderately polar compounds may actively play in the 5 α R inhibition.

Table 2. 5 α -reductase inhibitory activity of *T. indica* seed extracts.

Extract	Dutasteride equivalent 5 α R inhibitory activity (μg DE per mg of crude extract)
TIEE	50.11 ± 3.65
TIEAE	153.31 ± 10.60
TIHE	9.77 ± 3.82

Accurate mass screening was carried out using the UNIFI data analysis software. The base peak ions of distinct peaks were subjected to library matching using the Traditional Chinese Medicine Library. The base peak ion chromatograms of the TIEAE and blank are shown in Figures 2A and 2B, respectively. Blank peaks were discriminated from the chromatogram of the sample and were not annotated. Annotation of the candidate masses was based on the accurate mass match, isotopic ratio match, and precursor ion intensity counts. Peak annotations in the chromatogram are the retention time of the most intense peak in each spectrum. Each peak corresponds to one component sample. A total of 92 peaks were observed, 15 of which have putative identification based on the library search (Table 3). The 15 phytochemicals identified belonged to various classes such as phenolic compounds, tocopherols, fatty acids, phytosterol, saponins, and terpenoids.




Figure 2. Base peak ion chromatogram of TIEAE (A) and methanol, as blank (B).




Figure 2. (Cont.) Base peak ion chromatogram of TIEAE (A) and methanol, as blank (B).

Table 3. Putatively identified phytochemicals present in *T. indica* ethyl acetate extract based on the library search.

Component Name	Retention Time	Observed m/z	Error	Adducts
Catechin	3.02	291.0889	2.595	+H
Catechin-(4 α →8)-catechin	3.49	579.1483	-1.3544	+H, +Na, +K
Chrysin	4.51	255.0670	1.8247	+H
3-Hydroxy baicalein	4.83	287.0547	-0.2938	+H
Darutigenol	6.11	361.2051	-8.8839	+K
Nigakilactone I	7.83	399.1784	0.6304	+H, +Na, +K
24-Methylcholesta-7,22-dien-3 β -ol	8.40	399.3594	-2.7619	+H, +Na
Kushenol I	8.57	455.2041	-2.2984	+H
4,8,12-Trimethyltridecanoic acid	8.68	279.232		+Na
(20S)-3 α ,29-dihydroxylupan-27-oic acid	9.79	497.3618	1.6218	+Na, +K
Abrisapogenol B	9.99	497.3596	-0.4973	+Na
ar-Abietatriene	10.22	309.1947	-0.5379	+K
12-Hydroxyhydromethyl abietate	10.38	337.2724	-1.3166	+H
Vitamin E β	10.77	439.3529	-1.7232	+Na
3-O-[[β -D-Glucopyranosyl-(1→2)]- β -D-glucopyranosyl]-kaempferol	12.27	633.1488	6.1792	+H, +Na, +K

Studies have identified the use of *T. indica* in the management of BPH [18, 23]. In the present study, 5 α R which is important in the progression of BPH was found to be inhibited by TIEAE. The activity of *T. indica* against 5 α R may be attributed to the phytochemicals present. The seeds of *T. indica* are abundant in procyanidin [24], wherein the procyanidin B3 dimer, known as catechin-(4 α →8)-catechin [25] was putatively identified in TIEAE. In the study of Hiipakka et al. [26], procyanidin along with other phenolic compounds such as catechin, chrysin, baicalein, and kaempferol, possess various enzyme inhibitory effects against 5 α R.

The same study showed that the number and position of B-ring hydroxyl groups in phenolic compounds appear to be related to the varied bioactivity. The present study was also able to putatively identify Vitamin E β or β -tocopherol in TIEAE. In the molecular dynamic study of Khantam et al. [27], tocopherols including β -tocopherol were found to bind in multiple residues of 5 α R, affecting the enzyme activity. Long-chain saturated fatty acids also elicit inhibitory activity for 5 α R, with C12-C16 chains possessing enhanced inhibition [28]. Although no literature was found about the 5 α R inhibitory potential of 4,9,12-trimethyl-tridecanoic acid, its carbon chain number is within the 5 α R inhibitory criteria of long-chain saturated fatty acids reported by Liu et al. [28]. Furthermore, TIEAE was found to contain phytosterol, which may possess the potential to inhibit 5 α R [29, 30]. However, a high oral dose of phytosterol is needed to inhibit the enzyme as seen in the animal study of Awad et al. [31]. Lastly, terpenoids were also identified in TIEAE which may partly contribute to the bioactivity against 5 α R.

3. CONCLUSIONS

This study demonstrated the inhibitory potential of various extracts of *T. indica* seeds against 5 α R, TIEAE being the most active. Phenolic compounds, tocopherols, fatty acids, phytosterols, saponins, and terpenoids were putatively identified in the extract which may be the responsible phytochemicals in the enzyme inhibition. It is recommended, however, to conduct further studies on the specific phytochemicals putatively identified such as *in vivo* studies using animal models of BPH, to fully characterize its application in the urologic condition.

4. MATERIALS AND METHODS

4.1. Chemicals and reagents

Ethanol, ethyl acetate, and hexane used for extraction were of analytical grade (Ajax Finechem Pty Limited, New South Wales, Australia). Sucrose, testosterone, NADPH, NaOH, and Folin-Ciocalteu phenol were obtained from Sigma Aldrich (St. Louis, Missouri, USA) while dithiothreitol was procured from Promega (Madison, Wisconsin, USA). Solvents used for the HPLC-PDA analysis were of HPLC grade (Duksan, Ansan, South Korea), while solvents used for the ultra-performance liquid chromatography quadrupole time-of-flight mass spectrometry (UPLC-qTOF-MS) analysis were of LC-MS grade (Duksan, Ansan, South Korea), including the additive formic acid (Scharlab, Sentmenat, Spain). Additionally, type I Milli-Q[®] water (Millipore, Merck, Germany) was utilized. All other solvents and reagents used were of analytical grade and were obtained from local suppliers (Belman Laboratories, Quezon City, Philippines, and Harnwell Chemicals, Makati, Philippines).

4.2. *T. indica* crude seed extracts preparation

Fruits of *T. indica* were collected in the province of Abra in the Cordillera Administrative Region, Philippines. A sample was submitted to the Bureau of Plant Industry Manila under a voucher specimen number PLT-ID-CRPSD-2246-19 and was certified to be *T. indica* L. Fruit pods were manually cracked, and the seeds were separated from the fruit pulp. The seeds were washed, air-dried, and powdered using RT-12 pulverizing machine (Rong Tsong Precision Technology Co., Taichung, Taiwan). The resulting powdered *T. indica* seeds were separately macerated using three different solvents of varying polarities, namely, ethanol, ethyl acetate, and hexane. Maceration of seeds was performed for 48 h followed by 30 min sonication using Powersonic 410 sonicator (Human Lab Inc., Hwaseong, South Korea). The mixtures were filtered, and the filtrates were evaporated to dryness to obtain TIEE, TIEAE, and TIHE. The percent yield of the crude extracts was calculated. Afterward, they were placed in wide-mouth amber-colored bottles and stored under 2-8°C prior to the *in vitro* 5 α R inhibitory assay.

4.3. HPLC-PDA method

An HPLC-PDA method was used for the detection and quantification of testosterone in the *in vitro* 5 α R inhibitory assay. Waters Alliance e2695 HPLC system and 2998 PDA detector (Waters, Milford, Connecticut, USA) were used, equipped with HiQsil C18 column (4.6 x 150 mm, 5 μ m). Empower[®] 3 software was employed for data acquisition. Chromatographic conditions were optimized in terms of mobile phase composition, mode of elution, flow rate, UV detection, column temperature, and injection volume. For the selection of IS, triplicate runs were first conducted on testosterone with the candidate IS (i.e. propylparaben, ursodeoxycholic acid, and cholecalciferol) for their suitability in the method.

4.4. Mouse liver enzyme preparation

The 5 α R inhibitory activity of the three crude extracts was screened following the methods of Lee et al. [32] with modifications. Procedures for animal use were approved by the Institutional Animal Care and Use Committee of the University of the Philippines Manila (protocol number: 2019-034). Three healthy ICR mice were fasted overnight to decrease the concentration of liver glycogen and then anesthetized with 30 mg/kg of tiletamine-zolazepam (Zoletil[®], Virbac, Carros, France) followed by cervical dislocation. The livers were excised and washed with ice-cold homogenizing buffer (0.32 M sucrose, 0.001 M dithiothreitol, and 0.02 M potassium phosphate, pH 6.5). The livers were minced in a beaker with a pair of pre-sterilized scissors. The minced tissue was homogenized in the homogenizing buffer followed by centrifugation (MRC Laboratory-Instruments, Holon, Israel) at 3,940 xg for 10 min. The pellet produced was washed twice with 2 volumes of homogenizing buffer. The supernatants were combined from the two centrifugations and were centrifuged further at 3,940 xg for 1 h. The resulting pellet (microsome) was suspended in the homogenizing buffer and centrifuged at 3,940 xg for an additional 1 h. The microsome suspension was divided into small aliquots and was stored in an ultralow freezer (ThermoFisher Scientific, Asheville, North Carolina, USA) at -80°C prior to its use.

4.5. Microsomal protein content determination

The protein content of the microsome was determined using the modified Lowry method. The assay was carried out by diluting the sample to 1 ml with water and adding 0.9 ml of solution A (2 g/l KNaC₄H₄O₆ and 100 g/l Na₂CO₃ in 0.5 M NaOH) prior to a 10 min incubation at 50°C. The sample was cooled down to room temperature and 1 ml of solution B (0.2 g/l KNaC₄H₄O₆ and 0.1 g/l CuSO₄ in 0.1 M NaOH) was added. The mixture was left to stand for 10 min. Three ml of solution C (Folin-Ciocalteu phenol reagent in water, 1:16 v/v) was added and the mixture was incubated at 50°C for 10 min. A standard curve was made using bovine serum albumin, and the absorbance was read using a spectrophotometer (Hitachi High-Tech, Tokyo, Japan) at 600 nm [33].

4.6. Microsomal and sample solution preparation

The microsome was diluted immediately before use with 0.04 M potassium phosphate, pH 6.5, to achieve a 30 μ g protein concentration. The three crude extracts were dissolved separately in pure DMSO. From this stock solution, aliquots were diluted with 0.04 M potassium phosphate, pH 6.5, to obtain four concentrations (30, 60, 150, and 300 μ g/ml) of the sample solutions in 2% DMSO.

4.7. 5 α R assay

The method was based on the studies of Chaiyana et al. [29], Koseki et al. [34], and Morikawa et al. [35] with minor modifications in the determination of the 5 α R inhibitory activity of the extracts. Briefly, the reaction solution was preincubated without or with sample solution (50 μ l) in 0.001 M dithiothreitol (25 μ l), 1 mg/ml testosterone (75 μ l), and 1.54 mg/ml NADPH (125 μ l). The reaction was started with the addition of microsomal solution (250 μ l). The mixture was vortexed (Scientific Industries, Inc., Bohemia, New York, USA) for 3 s and was incubated for 10 min at 37°C. Afterward, the reaction was stopped by the addition of 1 ml ethyl acetate, followed by the addition of IS, 1 mg/ml propylparaben (125 μ l). Samples were vortexed for 60 s. The organic layer was collected and evaporated to dryness. The residue was dissolved in 3 ml methanol, passed through a 0.45 μ m syringe filter, and transferred to amber-colored HPLC vials for the quantification of testosterone using the HPLC-PDA method described above. Two additional reactions were prepared [36], namely, a solvent blank (sb) containing 2% DMSO instead of the sample solution and an enzyme blank (eb) that was mixed with ethyl acetate before the addition of the microsomal solution causing the non-conversion of testosterone to DHT. The % inhibition was calculated using the testosterone content defined in Eq. 2:

$$\text{(Eq. 2) \% inhibition} = [(T_{\text{ext}} - T_{\text{sb}}) / (T_{\text{eb}} - T_{\text{sb}})] \times 100$$

wherein T_{ext} , T_{sb} , and T_{eb} correspond to the unchanged testosterone levels in the sample, solvent blank, and enzyme blank solutions, respectively. Dutasteride, a well-known 5 α R inhibitor was used as the positive control. The IC₅₀ of dutasteride was calculated. Using the inhibitory equation of dutasteride, the DE values of extracts were calculated. Dilution factors for each concentration were considered. Triplicate measurements were performed. The extract with the highest inhibitory activity was subjected to phytochemical identification.

4.8. UPLC-qTOF-MS analysis

The extract with the highest inhibitory activity was subjected to phytochemical analysis using UPLC-qTOF-MS. The extract was dissolved in methanol, diluted to a final concentration of 0.5 mg/ml, and then

passed through a 0.2 μ m syringe filter. The analysis was performed on Waters Acquity UPLC Class I (Waters, Milford, Connecticut, USA). Five μ l of the sample were injected and separated on Acquity HSS T3 C18 column (2.1 x 100 mm, 1.8 μ m) at 40°C. The mobile phase was comprised of water + 0.1% formic acid (A) and acetonitrile + 0.1% formic acid (B). A 0.4 ml/min gradient was initiated. The Waters Xevo G2-XS qTOF, MS^E mode was run in a positive ion mode, with the capillary voltage set at 1.0 kV, cone voltage at 40 V, cone gas flow of 40 l/h, desolvation temperature at 550°C, desolvation gas flow of 950 l/h and source temperature at 120°C. The samples were run at a range of 50-1,200 m/z with a scan time of 0.15 s. Leucine-enkephalin was used as a reference for correction. Data were processed using the Waters UNIFI Scientific Information System v1.8.1.073 software and Waters Traditional Chinese Medicine Library.

4.9. Statistical analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) 27.0. The data gathered were recorded as mean \pm standard deviation (SD). Data were analyzed with one-way analysis of variance (one-way ANOVA) followed by Tukey's post hoc test. Mean values were considered to be statistically significant when $p < 0.05$.

Acknowledgement: The study was funded by the Philippine Institute of Traditional and Alternative Health Care.

Author contributions: Concept - G.L.T.D., R.A.M.M., E.C.A.; Design - G.L.T.D., R.A.M.M., E.C.A.; Supervision - E.C.A.; Resources - E.C.A.; Materials - E.C.A.; Data Collection and/or Processing - G.L.T.D., R.A.M.M.; Analysis and/or Interpretation - G.L.T.D., R.A.M.M.; Literature Search - G.L.T.D., R.A.M.M. Writing - G.L.T.D., R.A.M.M.; Critical Reviews - G.L.T.D., R.A.M.M., E.C.A.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Roehrborn CG. Pathology of benign prostatic hyperplasia. *Int J Impot Res.* 2008; Suppl 3(S11-18). <https://doi.org/10.1038/ijir.2008.55>
- [2] Lee C, Kozlowski JT, Grayhack JT. Intrinsic and extrinsic factors controlling benign prostatic growth. *Prostate.* 1997; 31(2): 131-138. [https://doi.org/10.1002/\(sici\)1097-0045\(19970501\)31:2%3C131::aid-pros9%3E3.0.co;2-q](https://doi.org/10.1002/(sici)1097-0045(19970501)31:2%3C131::aid-pros9%3E3.0.co;2-q)
- [3] Zitzmann M, Depenbusch M, Gromoll J, Nieschlag E. Prostate volume and growth in testosterone-substituted hypogonadal men are dependent on the CAG repeat polymorphism of the androgen receptor gene: a longitudinal pharmacogenetic study. *J Clin Endocrinol Metab.* 2003; 88(5): 2049-2054. <https://doi.org/10.1210/jc.2002-021947>
- [4] Lee CL, Kuo HC. Pathophysiology of benign prostate enlargement and lower urinary tract symptoms: current concepts. *Tzu Chi Med J.* 2017; 29(2): 79-83. https://doi.org/10.4103/tcmj.tcmj_20_17
- [5] Dehm SM, Tindall DJ. Molecular regulation of androgen action in prostate cancer. *J Cell Biochem.* 2006; 99(2): 333-344. <https://doi.org/10.1002/jcb.20794>
- [6] Azzouni F, Godoy A, Li Y, Mohler J. The 5 alpha-reductase isozyme family: a review of basic biology and their role in human diseases. *Adv Urol.* 2012; 530121. <https://doi.org/10.1155/2012/530121>
- [7] Chowdhury S, Beitel LK, Lumbroso R, Purisima EO, Paliouras M, Trifiro M. A targeted bivalent androgen receptor binding compound for prostate cancer therapy. *Horm Cancer.* 2019; 10(1): 24-25. <https://doi.org/10.1007/s12672-018-0353-6>
- [8] Wilson JD. The pathogenesis of benign prostatic hyperplasia. *Am J Med.* 1980; 68(5): 745-756. [https://doi.org/10.1016/0002-9343\(80\)90267-3](https://doi.org/10.1016/0002-9343(80)90267-3)
- [9] Imperato-McGinley J, Guerrero L, Gautier T, Peterson RE. Steroid 5alpha-reductase deficiency in man: an inherited form of male pseudohermaphroditism. *Science.* 1974; 184(4170): 1213-1215. <https://doi.org/10.1126/science.186.4170.1213>
- [10] Imperato-McGinley J, Zhu YS. Androgens and male physiology the syndrome of 5alpha-reductase-2 deficiency. *Mole Cell Endocrinol.* 2002; 198(1-2): 51-59. [https://doi.org/10.1016/s0303-7207\(02\)00368-4](https://doi.org/10.1016/s0303-7207(02)00368-4)
- [11] Russell DW, Wilson JD. Steroid 5 alpha-reductase: two genes/two enzymes. *Annu Rev Biochem.* 1994; 63: 25-61. <https://doi.org/10.1146/annurev.bi.63.070194.000325>
- [12] Carson C 3rd, Rittmaster R. The role of dihydrotestosterone in benign prostatic hyperplasia. *Urology.* 2003; 61(4 Suppl 1): 2-7. [https://doi.org/10.1016/s0090-4295\(03\)00045-1](https://doi.org/10.1016/s0090-4295(03)00045-1)
- [13] Edwards JL. Diagnosis and management of benign prostatic hyperplasia. *Am Fam Physician.* 2008; 77(10): 1403-1410.

- [14] Zaid H, Raiyn J, Nasser A, Saad B, Rayan A. Physicochemical properties of natural based products versus synthetic chemicals. *Open Nutraceutical J*. 2010; 3: 194-202. <http://doi.org/10.2174/18763960010030100194>
- [15] Boy HIA, Rutilla AJH, Santos KA, Ty AMT, Yu AI, Mahboob T, Tangpoong J, Nissapatorn V. Recommended medicinal plants as source of natural products. *Digital Chin Med*. 2018; 1(2): 131-142. [https://doi.org/10.1016/S2589-3777\(19\)30018-7](https://doi.org/10.1016/S2589-3777(19)30018-7)
- [16] Vadivel V, Pugalenthi M. Evaluation of nutritional value and protein quality of an under-utilized tribal food legume. *Indian J Tradit Knowl*. 2010. 9(4): 791-797.
- [17] Alok S, Jain SK, Verma A, Kumar M, Sabharwal M. Pathophysiology of kidney, gallbladder and urinary stones treatment with herbal and allopathic medicine: a review. *Asian Pac J Trop Dis*. 2013; 3(6): 496-504. [https://doi.org/10.1016/S2222-1808\(13\)60107-3](https://doi.org/10.1016/S2222-1808(13)60107-3)
- [18] Chika A, Bello SO, Yunusa A, Yahaya AB, Sadiq A. Ethnobotanical survey of medicinal plants used as remedies for benign prostatic hyperplasia in Sokoto North Western Nigeria. *Nat J Physiol Pharm Pharmacol*. 2022; 12: 511-518. <http://doi.org/10.5455/njppp.2022.12.01043202203032022>
- [19] Whorwood CB, Donovan SJ, Wood PJ, Phillips DIW. Regulation of glucocorticoid receptor α and β isoforms and type I 11 β -hydroxysteroid dehydrogenase expression in human skeletal muscle cells: a key role in the pathogenesis of insulin resistance? *J Clin Endocrinol Metab*. 86(5): 2296-2308. <https://doi.org/10.1210/jcem.86.5.7503>
- [20] Lakka NS, Kuppan C. Principles of chromatography method development. In: Boldura OM, Balta C, Awwad S. (Eds). *Biochemical Analysis Tools Methods for Bio-Molecules Studies*. IntechOpen Limited, United Kingdom, 2019. <https://doi.org/10.5772/intechopen.89501>
- [21] Zhu C, Goodall DM, Wren SAC. Elevated temperature HPLC: principles and applications to small molecules and biomolecules. *LCGC Asia Pacific*. 2005; 8: 48-59.
- [22] Maehre HK, Dalheim L, Edvisen GK, Elvevoll EO, Jensen IJ. Protein determination-method matters. *Foods*. 2018; 7: 5. <https://doi.org/10.3390/foods7010005>
- [23] Noumi E, Kolaipila B. Prostatic illnesses: epidemiology and treatment with medicinal plants in Maroua (Cameroon). *Int J Res Ayurveda Pharm*. 2011; 2(3): 822-829.
- [24] Sudjaroen Y, Haubner R, Wurtele G, Hull WE, Erben G, Spiegelhalder B, Changbumrung S, Bartsch H, Owen RW. Isolation and structure elucidation of phenolic antioxidants from tamarind (*Tamarindus indica* L.) seeds and pericarp. *Food Chem Toxicol*. 2005; 43(11): 1673-1682. <https://doi.org/10.1016/j.fct.2005.05.013>
- [25] Noreljaleel AEM, Wilhelm A, Bonnet SL. Analysis of commercial proanthocyanidins. Part 6: sulfation of flavan-3-ols catechin and epicatechin, and procyanidin B-3. *Molecules*. 2020; 25: 4980. <https://doi.org/10.3390/molecules25214980>
- [26] Hiipakka RA, Zhang HZ, Dai W, Dai Q, Liao S. Structure-activity relationships for inhibition of human 5 α -reductases by polyphenols. *Biochem Pharmacol*. 2002; 63(6): 1165-1176. [https://doi.org/10.1016/s0006-2952\(02\)00848-1](https://doi.org/10.1016/s0006-2952(02)00848-1)
- [27] Khantam C, Yooi W, Sringarm K, Sommano SR, Jiranusornkul S, Carmona FD, Nimlamool W, Jantrawut P, Rachtanapun P, Ruksiriwanich W. Effect on steroid 5- α reductase gene expression of Thai rice bran extracts and molecular dynamics study on SRD5A2. *Biology*. 2021; 10(4): 319. <https://doi.org/10.3390/biology10040319>
- [28] Liu J, Shimizu K, Kondo R. Anti-androgenic activity of fatty acids. *Chem Biodivers*. 2009; 6(4): 503-512. <https://doi.org/10.1002/cbdv.200800125>
- [29] Chaiyana W, Punyoyai C, Somwongin S, Leelapornpisid P, Ingkaninan K, Waranuch N, Srivilai J, Thitipramote N, Wisuitiprot W, Schuster R, Viernstein H, Mueller M. Inhibition of 5 α -reductase, IL-6 secretion, and oxidation process of *Equisetum debile* Roxb. ex Vaucher extract as functional food and nutraceutical ingredients. *Nutrients*. 2017; 9: 1105. <https://doi.org/10.3390/nu9101105>
- [30] Azizi A, Mumin NH, Shafqat N. Phytochemicals with anti 5- α -reductase activity: a prospective for prostate cancer treatment. *F100Res*. 2021; 10: 221. <https://doi.org/10.12688/f1000research.51066.3>
- [31] Awad AB, Hartati MS, Fink CS. Phytosterol feeding induces alteration in testosterone metabolism in rat tissues. *J Nutr Biochem*. 1998; 9(12): 712-717. [https://doi.org/10.1016/S0955-2863\(98\)00076-X](https://doi.org/10.1016/S0955-2863(98)00076-X)
- [32] Lee HH, Ho CT, Lin JK. Theaflavin-3,3'-digallate and penta-O-galloyl-beta-D-glucose inhibit rat liver microsomal 5 α -reductase activity and the expression of androgen receptor LNCaP prostate cancer cells. *Carcinogenesis*. 2004; 25(7): 1109-1118. <https://doi.org/10.1093/carcin/bgh106>
- [33] Hartree EF. Determination of protein: a modification of the Lowry method that gives a linear photometric response. *Anal Biochem*. 1972; 48(2): 422-427. [https://doi.org/10.1016/0003-2697\(72\)90094-2](https://doi.org/10.1016/0003-2697(72)90094-2)

- [34] Koseki J, Matsumoto T, Matsubara Y, Tsuchiya K, Mizuhara Y, Sekiguchi K, Nishimura H, Watanabe J, Kaneko A, Hattori T, Maemura K, Kase Y. Inhibition of rat 5 α -reductase activity and testosterone-induced sebum synthesis in hamster sebocytes by an extract of *Quercus acutissima* Cortex. Evid Based Complement Alternat Med. 2015; 853846. <https://doi.org/10.1155/2015/853846>
- [35] Morikawa T, Luo F, Manse Y, Sugita H, Saeki S, Chaiech S, Pongpiriyadacha Y, Muraoka O, Ninomiya K. Geranylated coumarins from Thai medicinal plant *Mammea siamensis* with testosterone 5 α -reductase inhibitory activity. Front Chem. 2020; 8: 199. <https://doi.org/10.3389/fchem.2020.00199>
- [36] Kumar T, Chaiyasut C, Rungseewijitprapa W, Suttajit M. Screening of steroid 5 α -reductase inhibitory activity and total phenolic content of Thai plants. J Med Plant Res. 2011; 5(7): 1265-1271.

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dspace.marmara.edu.tr>.