

# Comparison of the effect of UVB application on the antioxidant properties of different mushroom species with vitamins D<sub>2</sub> and D<sub>3</sub>

Aybike CEBECİ<sup>1\*</sup>, Turgut TAŞKIN<sup>2</sup>, Şule AKTAÇ<sup>3</sup>, Muhammet Emin ÇAM<sup>4</sup>, Rifat Sami SANCI<sup>5</sup>, Fatma Esra GÜNEŞ<sup>6</sup>

<sup>1</sup> Department of Nutrition and Dietetics, Institute of Health Sciences, Marmara University, İstanbul, Turkey.

<sup>2</sup> Department of Pharmacognosy, Faculty of Pharmacy, Marmara University, İstanbul, Turkey.

<sup>3</sup> Department of Nutrition and Dietetics, Faculty of Health Sciences, Marmara University, İstanbul, Turkey.

<sup>4</sup> Department of Pharmacology, Faculty of Pharmacy, İstanbul Kent University, İstanbul, Turkey.

<sup>5</sup> Elit Engineering, İstanbul, Turkey.

<sup>6</sup> Department of Nutrition and Dietetics, Faculty of Health Sciences, İstanbul Medeniyet University, İstanbul, Turkey.

\* Corresponding Author. E-mail: [aybike.cebeci@marmara.edu.tr](mailto:aybike.cebeci@marmara.edu.tr) (A.C.); Tel. +90-216-777 57 01.

Received: 29 April 2024 / Revised: 6 June 2024 / Accepted: 8 June 2024

**ABSTRACT:** This study aimed to determine the effect of UVB application on different mushrooms and their antioxidant properties and compare their properties with pure vitamins D<sub>2</sub> and D<sub>3</sub>. In the study, as without UVB, with medium UVB and with high UVB, respectively; 3 ethanol extracts each from White button (E1, E2 and E3), Shiitake (E4, E5 and E6) and King oyster mushroom (E7, E8 and E9) were prepared. The antioxidant activities of the samples were compared with pure vitamin D<sub>2</sub> and D<sub>3</sub> solutions using DPPH, FRAP, and CUPRAC methods. The results show that vitamins D<sub>2</sub> (0.277 mMFeSO<sub>4</sub>/mg extract) and D<sub>3</sub> (0.380 mMFeSO<sub>4</sub>/mg extract) had the highest FRAP values. In mushroom species, it was determined that E3 (0.216 mMFeSO<sub>4</sub>/mg extract) and E9 (0.273 mMFeSO<sub>4</sub>/mg extract) mushroom extracts showed higher activity. Similarly, it was detected that vitamins D<sub>2</sub> (0.609 mMrolox equivalent/mg extract) and D<sub>3</sub> (1.149 mMrolox equivalent/mg extract) had the highest CUPRAC values. In mushroom species, E3 (0.324 mMrolox equivalent/mg extract) and E9 (0.292 mMrolox equivalent/mg extract) extracts had the highest CUPRAC values. According to the DPPH experiment, E6 (IC<sub>50</sub>:0.386 mg/ml) and E9 (IC<sub>50</sub>:0.411mg/ml) extracts showed stronger free radical scavenging activity compared to other extracts, while vitamin D forms showed lower activity than mushroom extracts. Ultimately, it was concluded that the antioxidant properties of mushrooms were positively influenced by the application of UVB.

**KEYWORDS:** Mushrooms; antioxidant activity; ultraviolet rays; ergocalciferol; cholecalciferol.

## 1. INTRODUCTION

Mushrooms are regarded as a rich source of many nutrients, such as riboflavin, copper, selenium, vitamin D, potassium, chitin, dietary fibre, and β-glucan [1]. They also possess a multitude of secondary metabolites, encompassing polypeptides, steroids, terpenes, and phenolic compounds [2]. Particular attention is paid to phenolic compounds because of their potent antioxidant properties [3]. Vitamin D, which has two main forms, ergocalciferol (vitamin D<sub>2</sub>) and cholecalciferol (vitamin D<sub>3</sub>), is a fat-soluble vitamin that can be obtained through diet and is also a secosteroid prohormone produced in the skin by ultraviolet B (UVB) from sunlight [4,5]. While vitamin D<sub>3</sub> is synthesised from the 7-dehydrocholesterol in the skin via UVB radiation and obtained through animal-based foods, vitamin D<sub>2</sub> is produced by exposing mushroom ergosterol to UV rays [6,7].

Research on improving the antioxidant qualities of fresh and edible mushrooms shows that applying UVB radiation yields notable results in this area [8,9]. Similarly, various studies indicate that UV treated mushrooms are a potential plant-based source of dietary vitamin D<sub>2</sub>, due to their conversion of ergosterol to ergocalciferol [10,11]. Moreover, UVB light is considered safe and suitable for mushrooms because it enhances their nutrient profile and does not cause detrimental changes in their composition [12]. Therefore, enhancing both the nutritional profile and antioxidant capacity of mushrooms through this methodology may be regarded as an innovative strategy.

**How to cite this article:** Cebeci A, Taşkın T, Aktaç Ş, Çam ME, Sancı RS, Güneş FE. Comparison of the effect of UVB application on the antioxidant properties of different mushroom species with vitamins D<sub>2</sub> and D<sub>3</sub>. J Res Pharm. 2024; 28(5): 1508-1515.

Cells experience heightened oxidative stress, primarily due to the increased generation of reactive oxygen species (ROS). This stress happens when the body's antioxidant defence mechanisms exceed their ability to neutralise ROS formation. Excessive ROS levels can potentially damage cellular components, encompassing nucleic acids, lipids, and proteins [13]. Uncontrolled free radical production has been linked to more than 100 diseases, including atherosclerosis, diabetes, various cardiovascular diseases, cirrhosis, neurological disorders, cancer, lung diseases, drug-induced deafness, Parkinson's disease, etc. [14].

Faced with oxidative stress, the organism triggers diverse defence mechanisms to alleviate or counterbalance the detrimental effects of ROS. These responses are based on a rise in the activity of antioxidants and related enzymes [15]. By interacting with free radicals, antioxidants are compounds that can terminate chain reactions [16]. Hence, it is thought that the consumption of antioxidant-rich foods might contribute to alleviating the adverse consequences of oxidative damage [17].

In recent years, mushrooms have attracted attention in terms of the antioxidant compounds they contain. Mushrooms are also thought to provide a wider range of health benefits than pure vitamin D, as they are a natural food source, and both their antioxidant content and vitamin D<sub>2</sub> levels increase with exposure to UVB [8-11,18,19].

In this regard, the objective of this study is to evaluate how UVB application affects the antioxidant properties of different mushrooms and to compare their antioxidant properties with vitamins D<sub>2</sub> and D<sub>3</sub>.

## 2. RESULTS

The 2,2-diphenyl-1-picryl-hydrazyl (DPPH) experiment revealed that E6 (IC<sub>50</sub>:0.386 mg/ml) and E9 (IC<sub>50</sub>:0.411 mg/ml) mushroom extracts exhibited stronger activity to scavenge free radicals in comparison to the other extracts. It was also shown that vitamin D forms showed lower scavenging activity than other extracts except E7. All of the extracts, however, were found to have less capacity to scavenge free radicals than the standard ascorbic acid (Table 1).

Based on an analysis of the extracts' ferric ion reduction capabilities, vitamins D<sub>2</sub> (0.277 mMFeSO<sub>4</sub>/mg extract) and D<sub>3</sub> (0.380 mMFeSO<sub>4</sub>/mg extract) were shown to have the highest ferric reducing antioxidant power (FRAP) values. The ferric ion reduction capacity of E9 (0.273 mMFeSO<sub>4</sub>/mg extract) mushroom extract was found to be higher than that of other extracts in the category of mushrooms. Furthermore, as with the DPPH analysis, it was discovered that all extracts had lower levels of ferric ion-reducing activity than the reference compound, BHA (Table 1).

Vitamins D<sub>2</sub> (0.609 mMTE/mg extract) and D<sub>3</sub> (1.149 mMTE/mg extract) showed the greatest cupric ion reducing antioxidant capacity (CUPRAC) values when the Cu (II) reduction potentials of the extracts were assessed. It was discovered that the extracts E9 (0.292 mMTE/mg extract) and E3 (0.324 mMTE/mg extract) had a greater capacity to reduce Cu (II) in mushroom species than the other extracts. However, all samples had lower CUPRAC values compared to the standard compound (Table 1).

**Table 1.** Antioxidant activities of the samples

Extracts/standards	DPPH	FRAP	CUPRAC
	(IC <sub>50</sub> : mg/mL)	(mMFeSO <sub>4</sub> /mg extract)	(mMTE/mg extract)
	mean ± SD	mean ± SD	mean ± SD
E1	0.861 ± 0.013 <sup>d,e</sup>	0.107 ± 0.024 <sup>f</sup>	0.146 ± 0.009 <sup>e,f</sup>
E2	0.722 ± 0.016 <sup>c,d</sup>	0.199 ± 0.017 <sup>e</sup>	0.284 ± 0.002 <sup>d</sup>
E3	0.598 ± 0.05 <sup>b,c</sup>	0.216 ± 0.002 <sup>d,e</sup>	0.324 ± 0.102 <sup>d</sup>
E4	0.584 ± 0.121 <sup>b,c</sup>	0.100 ± 0.007 <sup>f</sup>	0.108 ± 0.037 <sup>f</sup>
E5	0.519 ± 0.027 <sup>b,c</sup>	0.105 ± 0.005 <sup>f</sup>	0.176 ± 0.007 <sup>e,f</sup>
E6	0.386 ± 0.043 <sup>b</sup>	0.110 ± 0.005 <sup>f</sup>	0.247 ± 0.075 <sup>d,e</sup>
E7	1.226 ± 0.090 <sup>f</sup>	0.109 ± 0.027 <sup>f</sup>	0.071 ± 0.009 <sup>f</sup>
E8	1.091 ± 0.228 <sup>e,f</sup>	0.260 ± 0.003 <sup>c,d</sup>	0.240 ± 0.003 <sup>d,e</sup>
E9	0.411 ± 0.109 <sup>b</sup>	0.273 ± 0.004 <sup>c</sup>	0.292 ± 0.017 <sup>d</sup>
D <sub>2</sub>	1.094 ± 0.068 <sup>e,f</sup>	0.277 ± 0.041 <sup>c</sup>	0.609 ± 0.025 <sup>c</sup>
D <sub>3</sub>	1.151 ± 0.048 <sup>f</sup>	0.380 ± 0.046 <sup>b</sup>	1.149 ± 0.063 <sup>b</sup>
BHA		5.81 ± 0.004 <sup>a</sup>	
Ascorbic acid	0.004±0.001 <sup>a</sup>		5.683±0.337 <sup>a</sup>

<sup>a</sup> Values are means ± standard deviation of triplicates.

<sup>b</sup> Values in a column with different superscripts are significantly different (p<0.05, Tukey).

<sup>c</sup> E1: White button mushroom with non-UVB, E2: White button mushroom with medium UVB, E3: White button mushroom with high UVB, E4: Shiitake mushroom with non-UVB, E5: Shiitake mushroom with medium UVB, E6: Shiitake mushroom with high UVB, E7: King oyster mushroom with non-UVB, E8: King oyster mushroom with medium UVB, E9: King oyster mushroom with high UVB, BHA: Butylated Hydroxyanisole

### 3. DISCUSSION

According to the main study outcomes, applying UVB to mushrooms significantly improved their antioxidant capacity.

Nowadays, mushrooms, recognised for their noteworthy biological properties and nutritional value, are a focal point of scientific inquiry, owing to their substantial influence on human health. The antioxidant capacities of mushrooms, particularly their ability to prevent free radical-induced cellular damage, are a major contributor to their health benefits [20].

#### 3.1. DPPH radical scavenging activity test

The assay on DPPH radical is useful in predicting antioxidant activities by inhibiting lipid oxidation. Therefore, the degree of free radical scavenging is determined by DPPH radical scavenging technique. The DPPH radical scavenging test is widely utilised because it produces rapid analytical results in a relatively brief period [21].

The IC<sub>50</sub> value of *Agaricus bisporus* ranges from 1.4 to 77.5 mg/ml, according to DPPH experiments on this mushroom offered in the scientific literature [22-24]. The IC<sub>50</sub> values of *Agaricus bisporus* with no UVB, medium UVB, and high UVB were reported to be 0.861, 0.722, and 0.598 mg/ml, respectively, in this investigation. Furthermore, these mushrooms demonstrated a superior capacity for scavenging DPPH radicals compared to results documented in the current literature.

Nam et al. reported that the IC<sub>50</sub> value of *Lentinula edodes* varied between 0.88-0.94 mg/ml, according to its growing environment [25]. In research examining the antioxidant properties of the water extract of *Lentinula edodes*, the IC<sub>50</sub> value of this mushroom was determined as 48.3 µg/ml [26]. The antioxidant properties of various *Lentinula edodes* sections (peel, fruiting body, gills, inner cap, and stipe) were investigated in a different study, and it was discovered that the samples' IC<sub>50</sub> values ranged from 0.129 to 0.213 mg/ml [27]. An investigation into how UVB exposure affects *Lentinula edodes*' antioxidant activity revealed that UVB treatment increased the mushroom's capacity to scavenge free radicals (DPPH) [9]. The IC<sub>50</sub> values of *Lentinula edodes* without UVB, with medium UVB, and with high UVB were determined to be 0.584, 0.519, and 0.386 mg/ml, respectively, in this search. Upon reviewing the literature, it was noted that there was inconsistency between the results obtained from the studies analyzing the IC<sub>50</sub> value of *Lentinula edodes*.

An investigation by Gąsecka and colleagues on the antioxidant characteristics of *Pleurotus eryngii* revealed an IC<sub>50</sub> value of 7.34 mg/ml [28]. IC<sub>50</sub> values of samples taken from different parts of *Pleurotus eryngii* (peel, fruiting body, gills, inner cap, and stipe) were observed to vary between 0.171 and 6.331 mg/ml in another study [27]. According to a study that assessed the antioxidant properties of excreted parts of *Pleurotus eryngii*, the IC<sub>50</sub> value of this mushroom ranged from 0.41 to 0.52 mg/ml [29]. IC<sub>50</sub> values of *Pleurotus eryngii* without UVB, with medium and high UVB, were 1.226, 1.091, and 0.411 mg/ml, respectively, in this investigation. As with *Lentinula edodes*, inconsistent results were obtained when the IC<sub>50</sub> levels of *Pleurotus eryngii* were examined by reviewing the literature.

#### 3.2. Ferric reducing antioxidant power (FRAP) assay

The FRAP assay, as an electron transfer-based method, measures the ferric ion ligand complex reduction by antioxidants in an acidic environment to the ferrous complex [30]. This assay assesses the reducing potential of a sample as a means to gauge its antioxidant capacity. Increased antioxidant activity is correlated with increased reduction potential [31].

*Agaricus bisporus* had FRAP values ranging from 1.35 to 11.62 µMTE/g in a study where Smolskaitė and colleagues tested the antioxidant properties of different mushroom species [32]. Gan et al., in their antioxidant analysis study on aqueous and 60% ethanol extracts of different mushrooms, determined the FRAP values of *Agaricus bisporus* as 186.72 and 84.69 µM Fe<sup>2+</sup> equivalent/g extract, respectively, according to these extraction types [23]. The FRAP values of *Agaricus bisporus* with no UVB, medium UVB, and high UVB were found to be 0.107, 0.199, and 0.216 mMFeSO<sub>4</sub>/mg extract, respectively, in this research. Reviewing the data, it became clear that, similar to other analyses, the FRAP values of *Agaricus bisporus* demonstrated variations in the literature.

Investigating the impact of heat and extraction duration on the antioxidant capacities of *Lentinula edodes*, Baptista et al. observed variations in the FRAP values of the mushroom samples, ranging from 0.434 to 1.197 mMTE/g of extract [33]. The FRAP value of *Lentinula edodes* was 64.79 mg/ml, according to research conducted by Carneiro et al. [34]. A study on how UVB rays affect the antioxidant properties of *Lentinula edodes* revealed that UVB application increased the FRAP value, parallel to this investigation [9].

The findings of the current study demonstrated that the FRAP values of *Lentinula edodes* without UVB, with medium UVB, and with high UVB were 0.100, 0.105, and 0.110 mMFeSO<sub>4</sub>/mg, respectively. It is imperative to underscore that FRAP values reported for *Lentinula edodes* show discrepancies across various studies in the literature.

During an examination of the antioxidant properties of *Pleurotus* species, the reported FRAP values for *Pleurotus eryngii* exhibited a range of 13.30 to 19.63 Fe<sup>2+</sup>/g extract levels [35]. In an alternative investigation, the FRAP value for *Pleurotus eryngii* was determined to be 18.25 mg TE/g extract [36]. The FRAP values of *Pleurotus eryngii* without UVB, with medium UVB, and with high UVB were evaluated as 0.109, 0.260, and 0.273 mMFeSO<sub>4</sub>/mg, respectively, in this experiment. Consistent with other analyses, the results identified in this study differ from those documented in the existing scientific literature.

### 3.3. The cupric ion reducing antioxidant capacity (CUPRAC) method

A copper reduction test called the CUPRAC assay was developed as an alternative to the FRAP test. In the CUPRAC test, copper acts as the oxidizing agent instead of iron. This procedure quantifies the potential of the antioxidants to convert cupric (Cu<sup>2+</sup>) ions to cuprous (Cu<sup>+</sup>) ions [30].

The CUPRAC values of *Agaricus bisporus* were from 0.07 to 1.19, according to a study searching the antioxidant activity of *Agaricus* species [37]. A study examining the bioactive characteristics of various mushroom species discovered that, depending on the extraction method, the CUPRAC value of *Agaricus bisporus* measured between 49.69 to 132.40 μMTE/g extract [38]. According to this study, *Agaricus bisporus* with no UVB, medium UVB, and high UVB had CUPRAC values of 0.146, 0.284, and 0.324 mM/mg extract, respectively. In comparison, it could be seen that, like other analyses, the results in the literature were in contradiction.

*Lentinula edodes* was reported to have CUPRAC values that differ from 8.04 to 19.51 mgTE/g extract in a study assessing its antioxidant activity [39]. In another investigation, the reported CUPRAC values for *Lentinula edodes* spanned from 0.155 to 2.578 mg/ml [40]. According to findings from a parallel investigation, the CUPRAC values of *Lentinula edodes* measured between 42.38 and 171.60 μMTE/g extract [32]. The current investigation yielded CUPRAC values of 0.108, 0.176, and 0.247 mMTE/mg extract for *Lentinula edodes* with no UVB, medium UVB, and high UVB, respectively. The outcomes were different when compared with relevant research in the literature.

*Pleurotus eryngii* was identified with CUPRAC values ranging from 0.125 to 2.657 mg/ml in a study exploring the antioxidant properties of different mushroom species [40]. The CUPRAC values of the three *Pleurotus eryngii* forms tested in this work were 0.071, 0.240, and 0.292 mMTE/mg extract for the forms with no UVB, medium UVB, and high UVB, respectively. Consistent with prior research, the findings of this study exhibited variances compared to the documented literature.

This observed situation is predicted to arise from a multitude of factors, such as variety among species, variations in cultivation methods, environmental conditions, extraction times and techniques, developmental phases during harvest, and different parts of the mushrooms having different components.

This study is envisioned to provide support to the current literature, considering the limited research available on the effects of UVB application on mushrooms and their antioxidant properties. Moreover, the absence of any literature documenting a concurrent examination of these mushroom species, coupled with a comparative analysis of their antioxidant properties against vitamins D<sub>2</sub> and D<sub>3</sub>, emphasizes the originality of this study and highlights its strength. Another notable aspect of this research stems from the lack of studies assessing vitamins D<sub>2</sub> and D<sub>3</sub> using DPPH, FRAP, and CUPRAC methods, which contributes to the distinctive strengths of the current study. However, a thorough review of the literature reveals a clear impact of extraction methods on antioxidant activity. Thus, the utilization of a single extraction method in this study imposes a limitation on the research. Furthermore, scientific literature illustrates the variability in antioxidant properties among different parts of mushrooms, so the fact that different parts of the mushroom were not examined in this study constitutes another limitation of the study.

## 4. CONCLUSION

Consequently, it was determined that UVB application positively affected the antioxidant properties of mushrooms in this study. However, to fully determine the effect of UVB on the antioxidant properties of mushrooms, further studies should be conducted on more mushroom species, using different extraction methods and applying UVB to different parts of the mushrooms. Additionally, future research could focus on the bioavailability of the enhanced nutrient profiles in mushrooms resulting from UVB application to better understand their potential health benefits.

## 5. MATERIALS AND METHODS

### 5.1. Supply of samples

White button (*Agaricus bisporus*), Shiitake (*Lentinula edodes*), and King oyster mushrooms (*Pleurotus eryngii*) with and without UVB application were obtained from the Elit Engineering Food Industry. Vitamins D<sub>2</sub> (SIGMA PHR1238) and D<sub>3</sub> (SIGMA BP787) were purchased from Sigma Aldrich.

### 5.2. Drying mushrooms and UVB application

The drying process was applied to non-UVB-treated mushrooms. The mushrooms were dried at 55 °C for approximately 15-17 hours by turning the trays 90° in the same direction every 2-3 hours and were pulverized with a homogenizer.

UVB-treated mushrooms were dried using the same method and sliced into 5-15 mm pieces. Then, UVB was applied to some of the three mushroom species at a medium level (200-222 mj/cm<sup>2</sup>) and to some at a high level (390-430 mj/cm<sup>2</sup>) for 90 seconds. After the process, the mushrooms were pulverized with a homogenizer.

### 5.3. Classification of mushrooms

Classification of mushrooms according to UVB application is shown in Table 2.

**Table 2.** Classification of mushrooms according to UVB application

UVB Application	Mushroom Species		
	<i>Agaricus bisporus</i>	<i>Lentinula edodes</i>	<i>Pleurotus Eryngii</i>
Non-UVB	E1	E4	E7
Medium UVB	E2	E5	E8
High UVB	E3	E6	E9

<sup>a</sup> E: Extract

<sup>b</sup> E1: White button mushroom with non-UVB, E2: White button mushroom with medium UVB, E3: White button mushroom with high UVB, E4: Shiitake mushroom with non-UVB, E5: Shiitake mushroom with medium UVB, E6: Shiitake mushroom with high UVB, E7: King oyster mushroom with non-UVB, E8: King oyster mushroom with medium UVB, E9: King oyster mushroom with high UVB

### 5.4. Extract preparation

Each sample weighed 10 grams and was mixed with 100 ml of ethanol. After waiting for 48 hours, it was filtered with coarse filter paper. After repeating this procedure three times, the resulting filtrate was evaporated in a steam bath to obtain a solid extract. Before usage, the extracts were kept under storage conditions at +4 °C.

### 5.5. In vitro antioxidant assays

The antioxidant properties of the mushrooms were compared with pure vitamin D<sub>2</sub> and D<sub>3</sub> solutions by FRAP assay, CUPRAC method, and DPPH radical scavenging test.

In the DPPH experiment, 10 µL each of the samples prepared at different concentrations (0.5-3 mg/mL) and the standard solution (ascorbic acid) prepared at the same concentration were taken, and 240 µL of 0.1 mM DPPH (in methanol) solution was added. The standard solution and the samples to which DPPH solution was added were vortexed for a minute and then kept at room conditions and in the dark for 30 minutes. Absorbances were measured against the reference at 517 nm in a microplate reader. Under the same circumstances, a control was prepared using 10 µL of methanol in place of the samples and standard substance. The average values were determined after the experiment was conducted three times [41].

FRAP assay was performed by modifying the Benzie and Strain method [42]. FRAP reagent was prepared and for 30 minutes kept at 37 °C. After mixing 190 µL of the FRAP reagent with 10 µL of either the standard [butylated hydroxy anisole (BHA)] or the sample (3 mg/mL), the increase in absorbance against the blank that was prepared by replacing the extract with methanol was measured at 593 nm in the 4th minute. FRAP values for the samples were given as mMFeSO<sub>4</sub>/mg extract.

The method developed by Apak et al. was used to compute the samples' CUPRAC values [43]. 60  $\mu\text{L}$  each of neocuproin, copper (II) and ammonium acetate buffer solutions were added into the microplate. Then, 60  $\mu\text{L}$  of samples at 0.5 mg/ml concentration and 10  $\mu\text{L}$  of methanol were added. For 60 minutes, the microplate was kept closed at room temperature. At 450 nm, the wavelength of absorption of the samples was measured against the reference. CUPRAC values for the samples were expressed as mMTE/mg extract.

The design of the study is shown in Figure.




Figure. The design of the study

## 5.6. Statistical analysis

Triplicate assays were performed to test the antioxidant capacity, and the findings were expressed as mean  $\pm$  standard deviations for all data. Using Minitab-18 software, the one-way ANOVA and post-hoc Tukey test were used to evaluate the statistical significance.  $P < 0.05$  was considered as statistically significant.

**Acknowledgements:** This research was financially supported by the Marmara University Scientific Research Committee (Project No: TDK-2022-10777) and Health Institutes of Türkiye (TUSEB) Group A Urgent R&D Project Call (Project No: 2022-ACIL-08-27971).

**Author contributions:** Concept - A.C., Ş.A., M.E.Ç., F.E.G.; Design - A.C., T.T., Ş.A., M.E.Ç., F.E.G.; Supervision - Ş.A., M.E.Ç.; Resources - T.T., Ş.A., M.E.Ç.; Materials - T.T., R.S.S.; Data Collection and/or Processing - A.C., T.T., R.S.S.; Analysis and/or Interpretation - A.C., T.T., Ş.A.; Literature Search - A.C., T.T.; Writing - A.C., Ş.A.; Critical Reviews - A.C., T.T., Ş.A., M.E.Ç., R.S.S., F.E.G.

**Conflict of interest statement:** The authors declared no conflict of interest for this publication.

## REFERENCES

- [1] Kalaras MD, Richie JP, Calcagnotto A, Beelman RB. Mushrooms: A rich source of the antioxidants ergothioneine and glutathione. *Food Chem.* 2017; 233: 429-433. <https://doi.org/10.1016/j.foodchem.2017.04.109>
- [2] Li L, Ng TB, Zhao L, Fu M, Chen RR, Jiang Y, Liu F. Correlation of antioxidant activity with content of phenolics in extracts from the culinary-medicinal abalone mushroom *Pleurotus abalones* Han, Chen et Cheng (*Agaricomycetidae*). *Int J Med Mushrooms.* 2005; 7(1&2): 237-242. <https://doi.org/10.1615/IntJMedMushr.v7.i12.220>.
- [3] Palacios I, Lozano M, Moro C, D'Arrigo M, Rostagno MA, Martínez JA, Garcia-Lafuente A, Guillamón E, Villares A. Antioxidant properties of phenolic compounds occurring in edible mushrooms. *Food Chem.* 2011; 128(3): 674-678. <https://doi.org/10.1016/j.foodchem.2011.03.085>
- [4] Jeon SM, Shin EA. (2018). Exploring vitamin D metabolism and function in cancer. *Exp Mol Med.* 2018; 50(4): 1-14. <https://doi.org/10.1038/s12276-018-0038-9>.
- [5] Gil Á, Plaza-Díaz J, Mesa MD. (2018). Vitamin D: classic and novel actions. *Ann Nutr Metab.* 2018; 72(2): 87-95. <https://doi.org/10.1159/000486536>.
- [6] Liu J. Vitamin D content of food and its contribution to vitamin D status: a brief overview and Australian focus. *Photochem Photobiol Sci.* 2012; 11(12): 1802-1807. <https://doi.org/10.1039/c2pp25150g>.

- [7] Pierret L, Suppa M, Gandini S, Del Marmol V, Gutermuth J. Overview on vitamin D and sunbed use. *J Eur Acad Dermatol Venereol*. 2019; 33: 28-33. <https://doi.org/10.1111/jdv.15316>.
- [8] Feng Y, Xu H, Sun Y, Xia R, Hou Z, Li Y, Wang Y, Pan S, Li L, Zhao C, Ren H, Xin G. Effect of light on quality of preharvest and postharvest edible mushrooms and its action mechanism: A review. *Trends Food Sci Technol*. 2023; 104119. <https://doi.org/10.1016/j.tifs.2023.104119>
- [9] Tiwari A, Singh G, Sharma V, Srivastava RK, Sharma S. Harnessing the potential of UVB irradiation for improving the nutraceutical properties of edible xylophilic mushroom dried powder. *LWT-Food Sci Technol*. 2021; 150: 111913. <https://doi.org/10.1016/j.lwt.2021.111913>
- [10] Leung MF, Cheung PCK. Vitamins D and D2 in Cultivated mushrooms under ultraviolet irradiation and their bioavailability in humans: A mini-review. *Int J Med Mushrooms*. 2021; 23(11): 1-15. <https://doi.org/10.1615/IntJMedMushrooms.2021040390>
- [11] Keegan RJ, Lu Z, Bogusz JM, Williams JE, Holick MF. Photobiology of vitamin D in mushrooms and its bioavailability in humans. *Dermatoendocrinol*. 2013; 5(1): 165-176. <https://doi.org/10.4161/derm.23321>
- [12] Simon RR, Phillips KM, Horst RL, Munro IC. Vitamin D mushrooms: comparison of the composition of button mushrooms (*Agaricus bisporus*) treated postharvest with UVB light or sunlight. *J Agric Food Chem*. 2011; 59(16): 8724-8732. <https://doi.org/10.1021/jf201255b>.
- [13] Ferreira IC, Barros L, Abreu R. Antioxidants in wild mushrooms. *Curr Med Chem*. 2009; 16(12): 1543-1560. <https://doi.org/10.2174/092986709787909587>
- [14] Khatua S, Paul S, Acharya K. Mushroom as the potential source of new generation of antioxidant: a review. *Res J Pharm Technol*. 2013; 6(5): 496-505.
- [15] Lushchak VI. Free radicals, reactive oxygen species, oxidative stress and its classification. *Chem Biol Interact*. 2014; 224: 164-175. <https://doi.org/10.1016/j.cbi.2014.10.016>
- [16] Kumar S. The importance of antioxidant and their role in pharmaceutical science-a review. *Asian J Res Chem Pharm Sci*. 2014; 1(1): 27-44.
- [17] Aziz MA, Diab AS, Mohammed AA. Antioxidant categories and mode of action. In: Shalaby E. (Ed). *Antioxidants*. IntechOpen Book Series, London, 2019, pp.3-22.
- [18] Sánchez C. Reactive oxygen species and antioxidant properties from mushrooms. *Synth Syst Biotechnol*. 2017; 2(1): 13-22. <https://doi.org/10.1016/j.synbio.2016.12.001>
- [19] Dal NE, İşlekel H. Vitamin D as an immunomodulatory and antioxidant molecule: association between vitamin D deficiency and systemic sclerosis. *Turk J Immunol*. 2019; 7(1): 57-68. <https://doi.org/10.25002/tji.2019.945>.
- [20] Valverde ME, Hernández-Pérez T, Paredes-López O. Edible mushrooms: improving human health and promoting quality life. *Int J Microbiol*. 2015: 376387. <https://doi.org/10.1155/2015/376387>
- [21] Baliyan S, Mukherjee R, Priyadarshini A, Vibhuti A, Gupta A, Pandey RP, Chang CM. Determination of antioxidants by DPPH radical scavenging activity and quantitative phytochemical analysis of *Ficus religiosa*. *Molecules*. 2022; 27(4): 1326. <https://doi.org/10.3390/molecules27041326>.
- [22] Gąsecka M, Magdziak Z, Siwulski M, Mleczek M. Profile of phenolic and organic acids, antioxidant properties and ergosterol content in cultivated and wild growing species of *Agaricus*. *Eur Food Res Technol*. 2018; 244: 259-268. <https://doi.org/10.1007/s00217-017-2952-9>
- [23] Gan CH, Nurul Amira B, Asmah R. Antioxidant analysis of different types of edible mushrooms (*Agaricus bisporus* and *Agaricus brasiliensis*). *Int Food Res J*. 2013; 20(3): 1095-1102.
- [24] Elmastas M, Isildak O, Turkekul I, Temur N. Determination of antioxidant activity and antioxidant compounds in wild edible mushrooms. *J Food Compos Anal*. 2007; 20(3-4): 337-345. <https://doi.org/10.1016/j.jfca.2006.07.003>
- [25] Nam M, Choi JY, Kim MS. Metabolic profiles, bioactive compounds, and antioxidant capacity in *Lentinula edodes* cultivated on log versus sawdust substrates. *Biomolecules*. 2021; 11(11): 1654. <https://doi.org/10.3390/biom11111654>
- [26] Elhousseiny SM, El-Mahdy TS, Awad MF, Elleboudy NS, Farag MM, Yassein MA, Aboshanab KM. Proteome analysis and in vitro antiviral, anticancer and antioxidant capacities of the aqueous extracts of *Lentinula edodes* and *Pleurotus ostreatus* edible mushrooms. *Molecules*. 2021; 26(15): 4623. <https://doi.org/10.3390/molecules26154623>
- [27] Krüzselyi D, Móric ÁM, Vetter J. Comparison of different morphological mushroom parts based on the antioxidant activity. *LWT-Food Sci Technol*. 2020; 127: 109436. <https://doi.org/10.1016/j.lwt.2020.109436>
- [28] Gąsecka M, Mleczek M, Siwulski M, Niedzielski P. Phenolic composition and antioxidant properties of *Pleurotus ostreatus* and *Pleurotus eryngii* enriched with selenium and zinc. *Eur Food Res Technol*. 2016; 242: 723-732. <https://doi.org/10.1007/s00217-015-2580-1>
- [29] He P, Li F, Huang L, Xue D, Liu W, Xu C. Chemical characterization and antioxidant activity of polysaccharide extract from spent mushroom substrate of *Pleurotus eryngii*. *J Taiwan Inst Chem*. 2016; 69: 48-53. <https://doi.org/10.1016/j.jtice.2016.10.017>
- [30] Shahidi F, Zhong Y. Measurement of antioxidant activity. *J Funct Foods*. 2015; 18: 757-781. <https://doi.org/10.1016/j.jff.2015.01.047>
- [31] Munteanu IG, Apetrei C. Analytical methods used in determining antioxidant activity: A review. *Int J Mol Sci*. 2021; 22(7): 3380. <https://doi.org/10.3390/ijms22073380>
- [32] Smolskaitė L, Venskutonis PR, Talou T. Comprehensive evaluation of antioxidant and antimicrobial properties of different mushroom species. *LWT*. 2015; 60(1): 462-471. <https://doi.org/10.1016/j.lwt.2014.08.007>

- [33] Baptista F, Campos J, Costa-Silva V, Pinto AR, Saavedra MJ, Ferreira LM, Rodrigues M, Barros AN. Nutraceutical potential of *Lentinula edodes*' spent mushroom substrate: A comprehensive study on phenolic composition, antioxidant activity, and antibacterial effects. *J Fungi*. 2023; 9(12): 1200. <https://doi.org/10.3390/jof9121200>
- [34] Carneiro AA, Ferreira IC, Dueñas M, Barros L, Da Silva R, Gomes E, Santos-Buelga C. Chemical composition and antioxidant activity of dried powder formulations of *Agaricus blazei* and *Lentinus edodes*. *Food Chem*. 2013; 138(4): 2168-2173. <https://doi.org/10.1016/j.foodchem.2012.12.036>
- [35] Leong CC, Ho WY, Yeap SK, Krishnen G, Chong ZX, Ho JS, Lim PT, Ten ST. (2021). Assessment of phylogenetic, growth, and antioxidant capacity of *Pleurotus* spp. in Malaysia. *J Food Process Preserv*. 2021; 45(6): e15554. <https://doi.org/10.1111/jfpp.15554>
- [36] Naknaen P, Itthisoponkul T, Charoenthaikij P. Proximate compositions, nonvolatile taste components and antioxidant capacities of some dried edible mushrooms collected from Thailand. *J Food Meas Charact*. 2015; 9: 259-268. <https://doi.org/10.1007/s11694-015-9231-x>
- [37] Öztürk M, Duru ME, Kivrak Ş, Mercan-Doğan N, Türkoglu A, Özler MA. In vitro antioxidant, anticholinesterase and antimicrobial activity studies on three *Agaricus* species with fatty acid compositions and iron contents: A comparative study on the three most edible mushrooms. *Food Chem Toxicol*. 2011; 49(6): 1353-1360. <https://doi.org/10.1016/j.fct.2011.03.019>
- [38] Alkin M, Söğüt E, Seydim AC. Determination of bioactive properties of different edible mushrooms from Turkey. *J Food Meas and Charact*. 2021; 15(4): 3608-3617. <https://doi.org/10.1007/s11694-021-00941-7>
- [39] Xiaokang W, Lyng JG, Brunton NP, Cody L, Jacquier JC, Harrison SM, Papoutsis K. Monitoring the effect of different microwave extraction parameters on the recovery of polyphenols from shiitake mushrooms: Comparison with hot-water and organic-solvent extractions. *Biotechnol Rep*. 2020; 27: e00504. <https://doi.org/10.1016/j.btre.2020.e00504>
- [40] Abdullah N, Ismail SM, Aminudin N, Shuib AS, Lau BF. Evaluation of selected culinary-medicinal mushrooms for antioxidant and ACE inhibitory activities. *eCAM*. 2012: 464238. <https://doi.org/10.1155/2012/464238>
- [41] Fu W, Chen J, Cai Y, Lei Y, Chen L, Pei L, Zhou D, Liang X, Ruan J. Antioxidant, free radical scavenging, anti-inflammatory and hepatoprotective potential of the extract from *Parathelypteris nipponica* (Franch. et Sav.) Ching. *J Ethnopharmacol*. 2010; 130(3): 521-528. <https://doi.org/10.1016/j.jep.2010.05.039>
- [42] Benzie IFF, Strain, JJ. Ferric Reducing/Antioxidant Power Assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol*. 1999; 299: 15-27. [http://dx.doi.org/10.1016/S0076-6879\(99\)99005-5](http://dx.doi.org/10.1016/S0076-6879(99)99005-5)
- [43] Apak R, Güçlü K, Özyürek M, Karademir SE. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J Agric Food Chem*. 2004; 52(26): 7970-7981. <https://doi.org/10.1021/jf048741x>

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dspace.marmara.edu.tr>.