

ÜRİNER SİSTEM TAŞI OLAN HASTLARDA İDRARLA SİTRAT ATILIMI

URINARY CITRATE EXCRETION IN PATIENTS WITH UROLITHIASIS

Tülay AKÇAY* – Dildar KONUKOĞLU** – Ali Rıza KURAL

SUMMARY

Urinary citrate excretion of of 31 healthy people and 67 patients with urolithiasis determined by colorimetric citrate lyase assay.

A statistically significant difference was detected between the healthy male (325 ± 75.9 mg/g Cr) and female (453 ± 132 mg/g Cr) ($p < 0.05$) and it was seen that mean urinary citrate excretion of patients (172.61 ± 89.27 mg/g Cr) was significantly lower than that of healthy people (391 ± 105.9 mg/g Cr) ($p < 0.001$).

There was not a significant difference between the urinary citrate excretions of primer (180.07 ± 90.5 mg/g Cr) and recurrent (172.2 ± 88.18 mg/g Cr) stone formers.

Therefore the decreasead citrate excretion is an important factor of stone formation.

Key words: Citrate, Urolithiasis, Stones.

ÖZET

İdrarla sitrat atılımı kolorimetrik sitrat liyaz yöntemiyle 31 sağlıklı ve 67 hastada belirlendi.

Sağlıklı kadın (453 ± 132 mg/g Cr) ile erkek (325 ± 75.9 mg/g Cr) arasında anlamlı fark saptandı ($p < 0.05$) ve hastaların idrarla sitrat atılımları (172.61 ± 89.27 mg/g Cr) sağlıklı kişilerinkinden (391 ± 105.9 mg/g Cr) anlamlı olarak düşük bulundu ($p < 0.001$).

Primer taşlı hastalarla (180.07 ± 90.5 mg/g Cr) tekrar eden taşlı hastaların (172.2 ± 88.18 mg/g Cr) idrarla sitrat atılımı arasında anlamlı bir fark saptanmamıştır.

Azalmış sitrat atılımı taş oluşumunda önemli bir faktördür.

* Department of Biochemistry, Cerrahpaşa Medical Faculty, University of Istanbul - TURKEY.

** Department of Urology, Cerrahpaşa Medical Faculty, University of Istanbul - TURKEY

INTRODUCTION

Urolithiasis is one of the ancient diseases. Until the last century, the incidence of bladder stones was very high and they were mainly composed of calcium oxalate or uric acid. In those days, the stones composed of calcium phosphate, magnesium ammonium phosphate and cystine were familiar are mostly seen in developing countries and in the industrialized countries. Urinary calculi composed of magnesium ammonium phosphate could be seen only in old men with prostatic obstruction. Simultaneously the incidence of renal calculi composed of calcium oxalate and calcium phosphate or calcium oxalate, gradually increases [1].

Urinary citrate is an important determinant for crystallization of calcium salts, and an inhibitor of calcium oxalate [2, 3] and calcium phosphate [4] crystal growth. It is frequently lower in calcium stone formers than in normal subjects [5; 6]. It is also well recognized that women, who usually excrete more citrate than men are less prone to develop calcium stones [6, 7, 8] and stone formation was 2 or 3 fold high in men than women [5]. It was, also shown that urinary citrate excretion in patients with urolithiasis was low than the normals but that their serum citrate level was equal [9].

Hypocitraturia was seen in all patients with urolithiasis, except in patients with primary hyperparathyroidism (PHPT) and hyperuricosuric patients with calcium oxalate stones [5].

The aim of this study, is to investigate urinary citrate excretion in patients with urolithiasis and in healthy people.

MATERIALS AND METHODS

The control group (male: 15, female: 16, average age: 45 ± 15) established from the healthy staff of our hospital after clinical, laboratory and radiological elimination of urinary stones and other diseases. Their creatinine clearances were, also, in normal range. The patients (male: 42, female: 25, average age: 53 ± 17) were selected from the patients with urinary stones admitted to Urology Department of our hospital.

The patients group, also, subdivided according to the recurrence of their disease and there were 42 patients with primary stones and 25 patients with recurrent stones.

All patients had one or more urinary stones. To determine whether hypocitraturia in patients group, caused by renal deterioration, creatinine clearances of all patients were, also detected and it was seen that they were all normal. The urinary calcium, magnesium and uric acid levels determined by Perkin Elmer model atomic absorption spectrophotometry and enzymatic uricase methods, respectively. At the time of collection all patients were free of urinary tract infection. None of the subjects had a special diet, except patients who were encouraged to maintain a high fluid intake.

24-hour urine samples were collected adding Penicilline G and streptomycine [10]. Citrate determination was performed with an enzymatic method modified by Welshman and Mc Cambridge [11].

RESULTS

The mean 24-hour urinary citrate excretion in control group was 391 ± 105.9 mg/g Cr and in male and female, it was 325 ± 75.9 mg/g Cr and 453 ± 132 mg/g Cr, respectively. When we compared these values of male and female, we detected a statistically significant difference ($p < 0.05$).

The mean 24-hour urinary citrate excretion in patients was 172.61 ± 89.27 mg/g Cr and in male and female, it was 164 ± 70 mg/g Cr and 190 ± 136 mg/g Cr, respectively. However, there was not any significant difference between the values of male and female patients. When we compared the values of patients with the values of healthy people, a significant difference could be determined.

The patients group also, subdivided according to the recurrence of stone formation and the values of patients with primary stone (180.07 ± 90.5 mg/g Cr) didn't show any significant difference than the values of patients with recurrent stones (172.2 ± 88.18 mg/g Cr).

Urinary citrates, calcium, magnesium and uric acid excretions of patients groups and control group were given in Table 1.

DISCUSSION

Urinary calculi is partly a result of physiologic and chemical deterioration of urine and anatomic defect [12]. The pathologic changes in urine are; high urinary calcium [13, 14], oxalate [15, 16] and uric acid [17] excretion, deficiency of protective inhibitors [18], as citrate [19] and super saturation of urine.

Table 1: The urinary citrate, calcium, magnesium, and uric acid excretions of controls and patients.

		Citrate (mg/g Cr)	Ca (mg/g Cr)	Mg (mg/g Cr)	Uric Acid (mg/g Cr)
Control	(n:31)	391.00±105.90	150.00±75.00	105.00±42.00	595.0±175
Patients	(n: 67)	172.61±89.27*	194.80±48.50	90.01±35.80	545.3±230
Primary Stones	(n: 42)	180.07±90.50*	205.90±45.20	95.45±40.50	525.5±255
Recurrent Stones	(n: 25)	172.20±88.18*	183.86±50.25	84.58±29.30	565.1±224

* Comparison with the control group $p < 0.001$

Citric acid is freely filtered in the renal glomerulus and is reabsorbed in the proximal renal tubule [20]. The presence of citrate in the urine is believed to be beneficial, because citrate chelates calcium, thereby reducing precipitation of calcium salts [21]. As calcium chelates with citrate, the concentration of free calcium ions decreases and therefore the supersaturation of urine needed for stone formation decreases.

Reduced urinary citrate excretion in stone formers has been documented in many publications [22, 23].

Some of the investigator reported that urinary citrate excretion was higher in women than in men [6, 7], which was also found in this study. The reason for the

higher citrate excretion in women couldn't been explained. However, it could not be possible to demonstrate a relationship between sex steroids and urinary citrate. Some of the investigators showed no difference between the urinary citrate excretion of men and women [24].

When all of these conflicting previous studies examined, it was seen that no protector added in urine specimens at the time of collection. As urine specimen usually contain bacteria fed with citrate as carbon source, the citrate content of urine specimens gradually decrease, till citrate determination performed. In our study, all urine specimens collected adding Penicilline G and streptomycine as these were known most effective protectors.

In this study, it was seen, urinary citrate excretion in patients with urolithiasis was significantly low than controls. However, there wasn't any difference between the levels of patients with primary and recurrent disease.

In conclusion, when our results and those of the preliminary studies are taken into consideration, low citrate excretion in patients with urolithiasis shows that the determination of urinary citrate excretion is very important.

REFERENCES

- 1- Landsdale K. Human stones. *Science*: 159, 1199-1207, 1968.
- 2- Robertson WG, Latif AB, Scurr RS, Sergenat V. Inhibitors of calcium oxalate crystallization in urine from stone formers. Abstract. Second International Urinary Stone Conference in Singapore, 1983.
- 3- Tiselius HG, Fornander AM. Evaluation of a routine method for determination of calcium oxalate growth inhibition in diluted urine samples. *Clin Chemistry* 27, 565-568, 1981.
- 4- Sutor DJ, Percival JM, Doonan S. Isolation and identification of some urinary inhibitors of calcium phosphate formations. *Clin Chim Acta*, 89, 273-278, 1978.
- 5- Hodgkinson A. Citric acid excretion in normal adults and in patients with renal calculus. *Clin Sci Mol Med*. 23: 203-212, 1982.

- 6- Tiselius HG. Urinar excretion of citrate in normal subjects and patients with urolithiasis. In: Smith LH, Robertson WG, Finlayson B, eds. Urolithiasis and basic research. New York: Pl Press 1981 pp: 39-44.
- 7- Nicar MJ, Skurla C, Sakhaee K, Pak CYC. Low urinary citrate excretion in nephrolithiasis. Urology 21: 8-14, 1983.
- 8- Hosking DH, Wilson JW, Liedtke RR, Smith LH, Wilson DM. Urinary citrate excretion in normal persons and patients with idiopathic calcium urolithiasis. J Lab Clin Med. 106: 682-689, 1985.
- 9- Minisola S, Rossi W, Bilgi F, Tabolli S, Mazzudi FG. Citric acid and calcium nephrolithiasis. Clin Chem. 34: 211-212, 1988.
- 10- Graef V, Schmidtman H, Jarrow K. On the preservation of urines for the determination of citrate. Abstracts of the 5th International Symposium on Urolithiasis and Related Clinical Research, 1984, p: 222.
- 11- Welschman SG, Mc Cambridge H. The estimation of citrate in serum and urine using a citrate lyase Technique. Clin Chim Acta, 46: 243-246, 1973.
- 12- Kok DJ, Papaoulos S, Bijuot OLM. Excessive crystal agglomeration with low citrate excretion in recurrent stone formers. Lancet 10, 1056-1058, 1986.
- 13- Hodgkinson A. Relations between oxalic, calcium magnesium and creatinine excretion in normal men and male patients with calcium oxalate kidney stones. Clin Sci. 46: 357-367, 1974.
- 14- Rudman D, Kutner MH, Reold SC, Waters WC, Gerron GG, Bleier J. Hypocitraturia in calcium nephrolithiasis. J Clin Endocrinol Metab. 55: 1052-7, 1982.
- 15- Marshall RW, Cochran M, Robertson WG, Hodgkinson A, Nordin BEC. The relation between the concentration of calcium salts in the urine and renal stone composition in patients with calcium-containing renal stones. Clin Sci. 43: 433-441, 1972.
- 16- Robertson WG, Peacock M, Nordin B. Activity products in stone forming and non-stone forming urine. Clin Sci. 34: 579-594, 1968.
- 17- Coe FL. Hyperuricosuric calcium oxalate nephrolithiasis. Kidney Int. 13: 418-426, 1978.
- 18- Dent CE, Sutor DJ. Presence or absence of inhibitor of calcium-oxalate crystal growth in urine of normals and stone-formers. Lancet. 2: 776-778, 1971.
- 19- Parks JH, Coe FL. A urinary calcium-citrate indeks for the evaluation of nephrolithiasis. Kidney International 30: 85-90, 1986.
- 20- Simpson DP. Citrate excretion: A window on renal metabolism. Am J Physiol 244: F223, 1983.
- 21- Baggio B, Gambaro G, Oliva et al. Calcium oxalate nephrolithiasis: An easy way to detect an imbalance between promoting and inhibiting factors. Clin Chim Acta 124: 149, 1982.
- 22- Nikkila M, Koivula T, Jokela H. Urinary citrate excretion in patients with urolithiasis and normal controls. Eur Urol 16: 382-385, 1989.
- 23- Akçay T, Konukoğlu D, Çelik Ç. Hypocitraturia in patients with urolithiasis. Arch. Dis. Child. 74: 350-351, 1996.
- 24- Grass L, Locke J, Rapoport A, Orcopoulos D. Colorimetric and enzymatic methods compared for assay citrate in urine. Clin Chem. 33: 1074, 1987.

(Received, January 10, 1995)