

The effects of N(1)-2,4-dihydroxybenzylidene-N-(4) hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV) on testicular damage in streptozotocin-induced diabetic rats

Sevim TUNALI¹ * , Tulay BAL-DEMIRCI² , Bahri ÜLKÜSEVEN^{2*} , Refiye YANARDAG¹ 

¹ Department of Chemistry, Biochemistry Division, Faculty of Engineering, Istanbul University-Cerrahpasa, Istanbul, Turkey.

² Department of Chemistry, Inorganic Chemistry Division, Faculty of Engineering, Istanbul University-Cerrahpasa, Istanbul, Turkey.

* Corresponding Author. E-mail: stunali@iuc.edu.tr (S.T.); Tel. +90-212-473 70 70/17791

Received: 23 September 2021 / Revised: 4 March 2022 / Accepted: 5 March 2022

ABSTRACT: Diabetes mellitus (DM) is a serious metabolic disorder that has negative effects on male sexual and reproductive functions in humans and animals. The purpose of current research is to demonstrate the effect of N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV) (VOL) on testicular damage in male rats with streptozotocin (STZ)-induced diabetes. Male Swiss albino rats were randomly grouped as follows: Control (intact) group animals; control group animals given VOL (0.2 mM/kg/day) for 12 days; STZ-induced diabetic animals; diabetic animals given VOL group, at same dose and time. Experimental diabetes was induced with a single dose of 65 mg/kg intraperitoneal STZ injection. On day 12, overnight fasted animals were sacrificed and testis tissues (right and left) were collected and homogenized in 0.9 % saline. After centrifugation, protein levels and non-enzymatic parameters such as glutathione, lipid peroxidation, protein carbonyl, as well as the activities of alkaline phosphatase, myeloperoxidase and enzymatic antioxidants were determined. Based on the results obtained, VOL was shown to be a potentially beneficial compound in the amelioration of damaged testicular tissue of male diabetic rats after 12 days of administration. Our results suggest that VOL may be a promising candidate for the development of new generation antidiabetic drugs, and its administration to diabetic rats may be a suitable candidate in reducing testicular damage.

KEYWORDS: Diabetes mellitus; oxovanadium complex; testicular damage; oxidative stress.

1. INTRODUCTION

Vanadium, an important transition element in glucose and lipid metabolism, is an insulin-mimetic, antilipidemic, and potent stress alleviating agent in Diabetes mellitus (DM) [1]. As knowledge about the biological importance and potentials of vanadium compounds becomes widespread, interest in the synthesis of new compounds and complexes of this element for therapeutic purposes is increasing [2-4]. Besides the medical and therapeutic activities of vanadium compounds, several coordination complexes including vanadyl (IV) ions have been studied in vitro and in vivo as insulin-mimetic agents [5].

Oxidative stress, caused by hyperglycemia, has been suggested to be an important factor in production of excessive reactive oxygen species (ROS). Disproportionately increased free radicals in diabetes by glucose oxidation, nonenzymatic glycation of proteins, and the subsequent oxidative degradation of glycated proteins declined antioxidant defense mechanisms and lead to damage of cellular organelles and enzymes [6]. This condition is associated with various pathological situations such as micro and macrovascular complications [7], nephropathy [8], cataract formation [9] and cardiovascular diseases [10].

Tunali S, Bal-Demirci T, Ülküseven B, Yanardag R. The effects of N(1)-2,4-dihydroxybenzylidene-N-(4) hydroxybenzylidene-S-methyl-thiosemicarbazidatooxovanadium(IV) on testicular damage in streptozotocin-induced diabetic rats. J Res Pharm. 2022; 26(3):675-686.

The developed complication of DM by oxidative stress consequently affect male reproductive functions [11]. In addition, researches on birth rates in developed countries have clearly demonstrated that increased DM incidence is intimately connected to falling birth and infertility rates [12,13]. Hyperglycemia decreases sperm motility and semen volume, it influences sperm counts and morphology, and as well as disrupt seminiferous tubular morphology [14,15]. The occurrence of male infertility in diabetes (characterized by hyperglycemia), is closely related to both increased production of ROS and impairment of the activity of testicular antioxidant enzymes [16,17]. Hence, oxidative stress in testicular tissue induce apoptosis of endothelium of germ cells, lipid peroxidation, and the oxidative damage of DNA and proteins [18,19]. Besides that, damage to the seminiferous tubules and serious decrease in sperm quality is observed in the reproductive organs of streptozotocin (STZ)-induced diabetic animals [20,21]. STZ, a diabetogenic agent widely used to induce experimental diabetes in animals, causes characteristic changes in blood insulin and glucose concentrations by acting on pancreatic beta cells [22].

Vanadium and vanadium compounds have been reported to mimic the action of insulin [5,7]. The most accepted opinion about this behavior as insulin-like agent is considered to be based on the protein tyrosine phosphatase (PTPs) inhibition [23].

In view of the fact that oxidative stress is known to be the one of the factors in the DM, we aim to scrutinize the effect of VOL (a newly developed compound) on ROS level and antioxidant capacity of the testicular tissue of STZ-induced diabetic animal model.

2. RESULTS

STZ-induced DM causes weight loss and muscle wasting occurs due to the increase in catabolytic reactions [5, 24]. The data from this set of animals, concerning body weights and fasting blood glucose levels have been previously published with respect to the synthesis, characterization and antidiabetic properties of VOL [5]. The body weight data were significant between all groups for 1, 6 and 12 days ($P_{ANOVA} = 0.001$; $P_{ANOVA} = 0.0001$; $P_{ANOVA} = 0.0001$ respectively). The loss seen in body weight of the diabetic animals was prevented by VOL administration which could be attributed to its anti diabetic effect (Table 1). Accordingly, significance was also observed between groups at day 1, 6 and day 12 for blood glucose levels ($P_{ANOVA} = 0.0001$; $P_{ANOVA} = 0.0001$; $P_{ANOVA} = 0.0001$ respectively). Blood glucose levels which were increased by STZ administration were diminished by VOL treatment (Table 2) [5].

Table 1. Mean levels of body weight for all groups (g)* (Yanardag et al., 2009) [5].

Groups	n	Day 0	Day 1	Day 6	Day 12	P _{t-test}
Control	5	247.81 ± 31.15	256.37 ± 32.22	259.10 ± 28.77	267.87 ± 33.81	0.801
Control + VOL	5	255.47 ± 17.29	262.71 ± 16.36	268.63 ± 14.82	279.08 ± 17.44	0.253
Diabetic	6	238.70 ± 26.37	215.16 ± 22.40	202.84 ± 26.88	194.01 ± 25.66	0.034
Diabetic + VOL	5	223.80 ± 5.41	203.42 ± 7.48	191.12 ± 23.25	187.37 ± 25.53	0.085
P_{ANOVA}		0.433	0.001	0.0001	0.0001	

*Data were presented as mean ± SD.

Abbreviations: n, number of animals; SD., standard deviation; ANOVA, analysis of variance; VOL, N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV); g., gram.

Table 2. Mean levels of blood glucose levels for all groups (mg/dl)* (Yanardag et al., 2009) [5].

Groups	n	Day 0	Day 1	Day 6	Day 12	P _{t-test}
Control	5	87.79 ± 2.90	90.58 ± 9.78	92.75 ± 7.82	91.80 ± 10.11	0.850
Control + VOL	5	83.64 ± 7.58	92.92 ± 8.62	90.64 ± 10.41	92.84 ± 8.03	0.322
Diabetic	6	84.60 ± 11.05	286.50 ± 25.78	307.99 ± 56.75	266.98 ± 78.83	0.0001
Diabetic + VOL	5	86.34 ± 9.50	251.89 ± 58.72	104.13 ± 21.61	99.17 ± 13.73	0.0001
P _{ANOVA}		0.893	0.0001	0.0001	0.0001	

*Data were presented as mean ± SD.

Abbreviations: n, number of animals; SD., standard deviation; ANOVA, analysis of variance; VOL, N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV).

With respect to testicular oxidative stress indices, lower level of glutathione (GSH) in STZ-induced diabetic group was observed ($p < 0.01$), whereas the levels of malondialdehyde (MDA) and protein carbonyl (PC) were elevated in comparison to intact rats ($p < 0.005$; $p < 0.005$). VOL administration to the diabetic animals significantly increased GSH level ($p < 0.05$) and significantly lowered the raised MDA and PC contents ($p < 0.05$; $p < 0.05$) (Table 3).

Table 4 represents the activities of ALP and MPO of all experimental groups. There was a significant alteration in the activities of alkaline phosphatase (ALP) and myeloperoxidase (MPO) in diabetic rats when compared to control group ($p < 0.01$; $p < 0.0001$). VOL treatment of diabetic group significantly prevented the increased activities of ALP and MPO resulting from diabetes ($p < 0.0001$; $p < 0.005$) (Table 4). Meanwhile, the activity of MPO in control animals given VOL complex significantly increased in comparison to intact control animals ($p < 0.0001$).

Table 3. Testis GSH, LPO and PC levels in control and experimental groups.

Groups	GSH (nmol GSH/mg protein)*	LPO (nmol MDA/mg protein)*	PC (nmol Carbonyl/mg protein)*
Control	0.98 ± 0.27	1.11 ± 0.24	0.34 ± 0.03
Control + VOL	0.71 ± 0.13	1.04 ± 0.27	0.32 ± 0.02
Diabetic	0.46 ± 0.12 ^a	1.82 ± 0.27 ^c	0.43 ± 0.02 ^c
Diabetic + VOL	0.86 ± 0.08 ^b	1.21 ± 0.39 ^b	0.36 ± 0.04 ^b
P _{ANOVA}	0.004	0.004	0.003

*Data were presented as mean ± SD.

^a $p < 0.01$ versus to control animals; ^b $p < 0.05$ versus to diabetic animals; ^c $p < 0.005$ versus to control animals.

Abbreviations: GSH, reduced glutathione; LPO, lipid peroxidation; PC, protein carbonyl; SD., standard deviation; ANOVA, analysis of variance; VOL, N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV).

Table 4. Testis ALP and MPO activities in control and experimental groups.

Groups	ALP (U/mg protein)*	MPO (U/g tissue)*
Control	41.94 ± 3.31	0.08 ± 0.01
Control + VOL	45.14 ± 4.17	0.23 ± 0.01 ^c
Diabetic	53.16 ± 4.52 ^a	0.15 ± 0.01 ^c
Diabetic + VOL	25.39 ± 2.40 ^b	0.10 ± 0.06 ^d
P_{ANOVA}	0.0001	0.0001

*Data were presented as mean ± SD.

^ap < 0.01 versus to control animals; ^bp < 0.0001 versus to diabetic animals; ^cp < 0.0001 versus to control animals; ^dp < 0.005 versus to diabetic animals.

Abbreviations: ALP, alkaline phosphatase; MPO, myeloperoxidase; SD., standard deviation; ANOVA, analysis of variance; VOL, N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV).

A significant change in enzymatic oxidative stress bio-markers of diabetic group when compared to control animals was observed (Table 5). A significant decrease in the activities of all enzymatic antioxidant enzymes in the STZ-induced hyperglycemic group as compared to the control animals testicular tissues was noticed ($p < 0.0001$; $p < 0.0001$; $p < 0.05$; $p < 0.0001$). On the other hand, VOL treatment to the diabetic rats significantly elevated the activities of catalase (CAT), glutathione peroxidase (GPx), glutathione-S-transferase (GST), and superoxide dismutase (SOD) ($p < 0.05$; $p < 0.0001$; $p < 0.0001$; $p < 0.05$).

Table 5. Testis CAT, GPx, GST and SOD activities in control and experimental groups.

Groups	CAT (U/mg protein)*	GPx (U/g protein)*	GST (U/mg protein)*	SOD (U/mg protein)*
Control	24.54 ± 0.34	0.40 ± 0.05	1.800 ± 0.02	15.18 ± 4.42
Control + VOL	18.05 ± 1.58 ^a	1.40 ± 0.11 ^a	0.150 ± 0.01 ^e	8.92 ± 1.24 ^a
Diabetic	19.63 ± 1.38 ^b	0.13 ± 0.03 ^b	0.153 ± 0.01 ^f	12.63 ± 3.34 ^b
Diabetic + VOL	22.51 ± 1.77 ^c	0.22 ± 0.02 ^d	0.206 ± 0.01 ^d	16.94 ± 1.15 ^c
P_{ANOVA}	0.0001	0.0001	0.0001	0.0001

*Data were presented as mean ± S.D.

^ap < 0.005 versus to control animals; ^bp < 0.0001 versus to control animals; ^cp < 0.05 versus to diabetic animals; ^dp < 0.0001 versus to diabetic animals; ^ep < 0.05 versus to control animals; ^fp < 0.05 versus to control animals.

Abbreviations: CAT; catalase; GPx, glutathione peroxidase; GST, glutathione-S-transferase; SOD, superoxide dismutase; SD., standard deviation; ANOVA, analysis of variance; VOL, N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium(IV).

In addition, a significant change in the activities of CAT, GPx, GST and SOD was observed in VOL administered control animals. The activities of CAT, GST and SOD of VOL treated control animals significantly decreased when compared to intact animals ($p < 0.005$; $p < 0.05$; $p < 0.005$). In the contrary to that, GPx activity increased significantly in VOL treated animals when compared to intact rats ($p < 0.005$) (Table 5).

3. DISCUSSION

DM, characterized by hyperglycemia, is a multifactorial disease. Nowadays, there is great concern that the increasing incidence of diabetes may affect the reproductive function of many men in active reproductive age [25]. The seen loss of body weight and increased blood glucose level of the diabetic animals were prevented by VOL, and this findings were discussed in detail in our previous research [5]. Numerous studies, both in vitro and in vivo, have demonstrated the insulin-mimetic properties of vanadium and its inorganic and organic compounds [7,8,26].

Experimental studies on vanadium compounds with different coordination geometries have shown that vanadium complexes with five coordination numbers are stronger inhibitors than those with coordination numbers six or seven [27]. In our previous researches and the current study, the new oxovanadium(IV) complex-VOL has been found to have a healing effect on many biochemical parameters in diabetic testicular tissue, as well as normalising hyperglycaemia [5,28,29]. This effect may in part be due to the inhibition of PTPs by vanadium in various ways. The PTPs active sites has a cysteine residue, and their catalytically mechanism is formed via fosfocysteine as an intermediate [23,27]. On the other hand, in the form of compound, vanadium taken into the cell binds to the active site of PTP. This is resulting with the inhibition of the enzyme and signal transduction pathways used in glucose uptake are activated [23, 30]. Vanadium mimics insulin in this way and causing the reduction of blood sugar concentration in the body [30].

Reduced glutathione is an endogenous tripeptide that plays an important role in inactivating hydrogen peroxide and lipid peroxides in the biological system. Reduced glutathione level is the marker of the oxidative status. Moreover, glutathione and its constituent cysteine are effective compounds for decarbonylation [31]. Thiols cause reduction in the levels of protein carbonyl content, since they affect protein conformation, activity and function [32]. In the current study, vanadium complex caused a rise in the level of GSH, while PC content was significantly decreased. The opposite of these was observed in the hyperglycemic group, GSH concentration decreased in the diabetic group as compared to the intact group, while a significant increase in the PC level was observed. Similar to our findings, the decreased level of GSH in the diabetic testicular tissue was also reported by Bal et al., (2011) [33]. This may be as a result of the cleavage of GSH into cysteine, a decline in GSH synthesis, or a decrease in cellular antioxidant reserves as a result of reactions such as mixed disulfide formation [34].

Sperm cells contain high concentrations of specific lipid compounds such as plasmalogen, sphingomyelin and polyunsaturated fatty acids. Insufficiency of antioxidant mechanisms makes polyunsaturated fatty acids more vulnerable to oxidative damage by the excessively produced ROS species in DM and causes damage to testicular tissue [35,36]. Furthermore, the negative effects of LPO in membrane structure can influence the ability of spermatozoa to take part in membrane fusion events that are relevant for fertilization. Reports provides evidence that defective sperm function is commonly induced by oxidative stress. Thus, affecting sperm motility via lipid peroxidation and altered DNA integrity [37]. In the present investigation, there is show significantly high LPO level in diabetic testicular tissue. Related finding like ours has also been reported in the study with diabetic testicular damage ameliorated by diosgenin [38]. Oral administration of VOL to diabetic rats caused a significant reduction of LPO level. This beneficial effect of the vanadium complex may be due to its hypoglycemic effect.

Vanadium and its compounds have been found to have an inhibitory effect on PTPs as well as enzymes that especially catalyze phosphate reactions [39]. Various phosphatases such as acid phosphatase, alkaline phosphatase and tyrosine-protein phosphatases can be inhibited by vanadium compounds. Alkaline phosphatases are membrane bond glycoprotein enzymes which catalyse the hydrolysis of phosphate monoesters from molecules. They also plays a role in the transfer of phosphate groups to hydroxyl groups of organic molecules. The catalytic mechanism of such enzymes is the phosphorylation of a serine residue on

the active side and subsequent transfer of the phosphate group to organic acceptor or water molecule [40]. The reason why vanadium and its complexes inhibit such enzymes is due to the physicochemical similarity of phosphate and vanadate, and their participation in similar reactions. Despite the structural similarity, it is important to emphasize that coordination geometries of vanadium complexes shows a greater flexibility. Besides that, as mentioned before, this strong inhibition can be due to five numbered coordination geometry of vanadium complexes [27,40]. In the current study, the new oxovanadium complex-VOL demonstrates a five coordination number. The inhibition of ALP in the diabetic group given VOL may therefore suggest that VOL has a role in protecting testicular tissue against STZ-induced diabetic oxidative injury.

MPO is the enzyme that catalyzes the production of reactive oxygen species including hypochlorous acid (HOCl) in the presence of hydrogen peroxide and halides. The findings of Aratani (2018), obtained from real-time PCR techniques showed that the rat's testicles and bulbourethral glands express the MPO gene [41,42]. Further, a different research observed that MPO activity is significantly higher in the testicular tissue of the rats with non-insulin dependent diabetes mellitus (NIDDM), thereby revealing the association of the MPO gene with T2DM [43,44]. Khosravi et al., (2019) also demonstrated an increased activity of MPO (a biomarker and consistent indicator of neutrophil proliferation and occurrence of inflammation) in diabetic animal testis [38]. In agreement with these findings, we observed an increased MPO activity in the diabetic animals testicular tissues. VOL administration which is associated with lower inflammation was effective in the improvement of MPO activity in testicular tissues of diabetic rats.

The pathological state in DM is strictly associated with an imbalance of ROS production and their effect on enzymatic antioxidant defense system in tissues [45]. The seminal vesicles have been shown to secrete CAT, GPx, GR, SOD, and GSH into the seminal fluid, as an additional antioxidative support for the spermatozoa [46,47]. Consistent with our findings on enzymatic antioxidant activities in diabetic testicular rat tissue, there is reported the decreased activity of antioxidant enzymes CAT, GPx and SOD [38]. The mechanism responsible for the protective effect of vanadium on testicular function may involve the enhancement of intracellular GSH level and increased activities of CAT, GPx, GST and SOD [46,48]. Additionally, the probability that vanadium mimics the insulin is partially owing to changes in prooxidant/oxidant balance [49,50]. The competitive behavior of vanadate with respect to phosphate is likely an indication of its insulin-mimetic or insulin-enhancing effect of vanadium compounds. In this way, an enhancement of the antioxidant defense of the testis may counteract the effects of the excessive oxidative stress.

A remarkable finding in our study was the observation that VOL administration to control animals resulted in reduction of the activities of testicular CAT, GST and SOD. Whereas, the activities of MPO and GPx significantly increased. The changes noticed upon vanadium treatment may be as a result of increased formation of oxygen-free radicals, which in turn overwhelm the body's antioxidant defense system. Findings of the present study showing altered enzymatic activities suggests that administration of vanadium complex may cause toxic effects in normal testicular tissue metabolism. Similar observations have been shown in different studies of vanadium toxicity [51], however a more comprehensive study is needed to be conducted to assess the safe and effective doses of VOL.

4. CONCLUSION

Modern pharmacological research suggests that testicular damage are caused by hyperglycemic-induced oxidative stress, apoptosis, and disorders associated with endocrine metabolism. Therefore, researchers have recently tried to control the development of hyperglycemic related reproductive damage by focusing on the synthesis of new chemical compounds that may mimic and replace insulin. The present study provided information on the effects of VOL, a novel vanadium complex, on diabetic testicular tissue, which can be presented as a possible therapeutic alternative in reducing diabetes-related male infertility.

5. MATERIALS AND METHODS

5.1. Synthesis of VOL

The oxovanadium(IV) complex (VOL) (Figure 1) was freshly prepared by starting from S-methyl-isothiosemicarbazide as earlier described [5]. In first step, 2-hydroxybenzaldehyde S-methyl-isothiosemicarbazone was obtained from reaction of S-methyl-isothiosemicarbazide (0.460 g) and 2,4-dihydroxybenzaldehyde (0.604 g). Then, the thiosemicarbazone (1mmol) and 2-hydroxybenzaldehyde (1 mmol) were dissolved in ethanol (50 mL). Vanadyl sulphate ($\text{VO}\text{SO}_4 \cdot 5\text{H}_2\text{O}$, 1 mmol) dissolved in 25 mL alcohol was added to the mixture. After stirring for 5 hours, the brownish-powder formed was filtered. The crude product was recrystallized from alcohol-ether (1:1) and checked its purity by TLC. The chemical structure of VOL was confirmed using elemental analysis and infrared spectrum. A yield of 64%, and m.p. > 380°C was obtained. For $\text{C}_{16}\text{H}_{13}\text{N}_3\text{O}_4\text{SV}$ (calc.): C, 48.82 (48.74); H, 3.24 (3.32); N, 10.59 (10.64); S, 8.23 (8.14). IR (ATR, cm^{-1}): (OH) 3410, (C=N) 1606, 1594, 1574, (CO) 1146, (V=O) 984, (VO) 478-436.

5.2. Experimental animals and study design

Animals: The experiments were reviewed and approved by Animal Care and Use Institute's Committee of the Istanbul University. Twenty-one male Swiss albino strain rats (200 ± 50 g) were kept in normal temperature ($22 \pm 2^\circ\text{C}$), and fed with standard chow and water ad libitum.

Inducement of diabetes: Diabetes was induced by a single intraperitoneal dose of streptozotocin (STZ) (65 mg/kg body weight) dissolved in cold citrate buffer (0.1 M, pH 4.5) [52]. After 24 hour of STZ injection, DM was confirmed by measuring fasting blood glucose levels from the rats tail. Rats with a fasting blood glucose level above 250 mg/dL were considered as diabetic. The part of the study related to weight and blood glucose values was detailed in our previous research [5].

Experimental Protocol: The rats were randomly divided into four groups as follows: two control groups divided as (1) control group-intact (n=5) and (2) treated control-received VOL (n=5) (0.2 mM/kg/day); two diabetic groups separated as (3) diabetic control (n=6) (65 mg/kg body weight) and (4) VOL treated diabetic group (n=5) (receiving the same doses of STZ and VOL). On the last day of the experiment (day 12), the mice were sacrificed under anesthesia.

5.3. Biochemical assays

Both testicular tissues, right and left, were homogenized in a cold saline solution and centrifuged at $10,000 \times g$ for 10 minutes. The obtained clear upper phase was used for estimation of biochemical parameters.

Total protein determination: Testis total protein level was determined by the method of Lowry [53]. Briefly, alkaline proteins are reacted with copper ions and then reduced by Folin reactive. The absorbance of the product was evaluated at 500 nm by a spectrophotometer and calculated to express the results of the parameters per protein.




Figure 1. The VOL, a thiosemicarbazone-based oxovanadium(IV) complex.

Glutathione determination: GSH level was quantified by measuring the content of -SH groups via spectrophotometric technique according to the Beutler (1971) at 412 nm [54]. Metaphosphoric acid was used for protein precipitation, and 5,5'-dithiobis-2-nitrobenzoic acid was used for color development. The results were expressed as nmol GSH/mg protein.

Lipid peroxidation determination: Testis malondialdehyde (MDA) level, an end product of lipid peroxidation (LPO), was determined as thiobarbituric acid reactive substances by the method of Ledwozyw (1986) by measuring the formation of malondialdehyde (MDA) [55]. LPO was expressed in terms of MDA equivalents as nmol MDA/mg protein.

PC determination: Testis tissue PC was determined spectrophotometrically by the method of Levine et al. (1990) [56]. Testis homogenate samples (0.5 ml) were incubated in dark at room temperature with 2 ml DNPH (10 mM) for 1 h and mixed every 15 min. About 2.5 ml 20% trichloroacetic acetic (TCA) were added and samples were kept in ice for 5 min. After centrifugation, the supernatants were discarded and precipitates were washed three times with a mixture of 2 ml of ethanol and ethyl acetate (1:1). The final precipitates were treated with 1 ml of guanidine hydrochloride (6 M) and incubated at 37°C for 10 min. The absorbances were read at 370 nm, and the results expressed in nmol carbonyl/mg protein.

ALP determination: The specific activities of alkaline phosphatase (ALP) were determined by the method of Walter and Schutt (1974) [57]. Under optimum conditions for measurements of the ALP, the absorbance was monitored at 405 nm. Enzyme activities were expressed as U/mg protein.

MPO determination: Enzymatic activity of MPO was determined by the method of Wei and Frenkel (1991) [58]. This involves a solution of tissue homogenate, phenol, H₂O₂, and 4-aminoantipyrine as color-generating substance. One unit of enzyme activity was defined as the amount of the MPO present per gram of protein which caused a change in absorbance per minute at 460 nm and 37°C. The results were expressed as U/mg protein.

CAT determination: Testis catalase (CAT) activity was determined by the method of Aebi [59]. The method is based on converting hydrogen peroxide (H₂O₂) to water by the effect of CAT. The absorbance of the samples was measured in 240 nm by a spectrophotometer. The results were expressed as U/mg protein.

GPx determination: GPx activity of testis tissues was determined by the method described by Paglia and Valentine (1967) and modified by Wendel (1981) [60,61]. GPx catalyzes the conversion of hydrogen peroxide (H₂O₂) to water. The generated glutathione disulfide (GSSG) is reduced to GSH with consumption of nicotinamide adenine dinucleotide phosphate (NADPH) by glutathione reductase. During the oxidation of NADPH to NADP, the decrease in absorbance was measured spectrophotometrically at a wavelength of 366 nm. GPx activity was calculated using an extinction coefficient of 6.22 mM⁻¹ cm⁻¹, and the results are expressed as U/g protein.

GST determination: GST is a metabolizing enzyme that plays a prominent role in the detoxification of oxidized metabolites and may serve as an antioxidant. GST enzyme activity was measured according to the method of Habig and Jacoby (1981) [62]. GST catalyzes the reaction between GSH and 1-chloro-2,4-dinitrobenzene. The resultant product formed was monitored spectrophotometrically at 340 nm and 25°C. The level of GST activities in testicular tissue was evaluated by an extinction coefficient of 0.625 mM⁻¹ cm⁻¹, and expressed as U/mg protein.

SOD determination: SOD activity was assayed according to the method described by Mylroie, Collins, Umbles, and Kyle (1986) [63]. The reaction mixture contains potassium phosphate buffer (pH 7.8), o-dianisidine dihydrochloride, and riboflavin. The reaction was initiated by the addition of riboflavin to the reaction mixture and the absorbance of the mixture was monitored immediately at a wavelength of 460 nm. Enzyme activity was determined using the SOD standard and activities in testicular tissue were expressed as U/mg protein.

The analysis of each sample was made in duplicate.

5.4. Statistical analyses

Results were evaluated using an unpaired t test and analysis of variance (ANOVA) using the NCSS statistical computer package. The values were expressed as mean ± SD. p < 0.05 was considered as significant.

Acknowledgements: This study was supported by Research Fund of Istanbul University Project Number: UDP-

3966/24062009.

Author contributions: Concept – S.T., T.B.D., B.U., R.Y.; Design – S.T., T.B.D., B.U., R.Y.; Supervision – B.U., R.Y.; Resources – S.T., T.B.D., B.U., R.Y.; Materials – S.T., T.B.D.; Data Collection and/or Processing – S.T., T.B.D., B.U., R.Y.; Analysis and/or Interpretation – S.T., T.B.D.; Literature Search – S.T., T.B.D., B.U., R.Y.; Writing – S.T., T.B.D., B.U., R.Y.; Critical Reviews – B.U., R.Y.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Tripathi D, Mani V, Pal RP. Vanadium in biosprocesses. *Biol Trace Elem Res.* 2018; 186(1): 52-67. [CrossRef]
- [2] Naglah AM, Al-Omar MA, Almehezia AA, Obaidullah AJ, Bhat MA, Kalmouch A, Al-Wasidi AS, Al-Humaidi JY, Refat MS. Synthesis, characterization, and anti-diabetic activity of some novel vanadium-folate-amino acid materials. *Biomolecules.* 2020; 10(5): 781.[CrossRef]
- [3] Ceylan BI. Oxovanadium (IV) and nickel (II) complexes obtained from 2,2'-dihydroxybenzophenone-S-methyl-thiosemicarbazone: Synthesis, characterization, electrochemistry, and antioxidant capability. *Inorg Chim Acta.* 2021; 120186. [CrossRef]
- [4] Ceylan BI. Oxovanadium (IV)-containing N2O2 chelate complex; crystal structure determination and DFT. *JTCSCA.* 2016; 3(3): 393-402. [CrossRef]
- [5] Yanardag R, Demirci TB, Ulküseven B, Bolkent S, Tunali S, Bolkent S. Synthesis, characterization and antidiabetic properties of N(1)-2,4-dihydroxybenzylidene-N(4)-2-hydroxybenzylidene-S-methyl-thiosemicarbazidato-oxovanadium (IV). *Eur J Med Chem.* 2009; 44(2): 818-826. [CrossRef]
- [6] Maritim AC, Sanders RA, Watkins JB 3rd. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol.* 2003;17(1):24-38. doi: 10.1002/jbt.10058. PMID: 12616644.
- [7] Akgün-Dar K, Bolkent S, Yanardag R, Tunali S. Vanadyl sulfate protects against streptozotocin-induced morphological and biochemical changes in rat aorta. *Cell Biochem Funct.* 2007;25(6):603-609. [CrossRef]
- [8] Yanardag R, Bolkent S, Karabulut-Bulan O, Tunali S. Effects of vanadyl sulfate on kidney in experimental diabetes. *Biol Trace Elem Res.* 2003; 95(1): 73-85. [CrossRef]
- [9] Feldman-Billard S, Dupas B. Eye disorders other than diabetic retinopathy in patients with diabetes. *Diabetes Metab.* (2021); 101279. [CrossRef]
- [10] Bayrak BB, Koroglu P, Karabulut Bulan O, Yanardag R. Metformin protects against diabetes-induced heart injury and dunning prostate cancer model. *Hum Exp Toxicol.* 2021; 40(2): 297-309. [CrossRef]
- [11] Alabi TD, de Villiers C, du Plessis SS, Monsees TK, Brooks NL, Oguntibeju OO. The beneficial role of anchomanes difformis in STZ-induced reproductive dysfunction in male wistar rats. *Diabetes Metab Syndr Obes.* 2020; 13: 4543-4560. [CrossRef]
- [12] Shi GJ, Li ZM, Zheng J, Chen J, Han XX, Wu J, Li GY, Chang Q, Li YX, Yu JQ. Diabetes associated with male reproductive system damages: Onset of presentation, pathophysiological mechanisms and drug intervention. *Biomed Pharmacother.* 2017; 90: 562-574. [CrossRef]
- [13] Bahmanzadeh M, Vahidinia A, Mehdinejadi S, Shokri S, Alizadeh Z. Dietary supplementation with astaxanthin may ameliorate sperm parameters and DNA integrity in streptozotocin-induced diabetic rats. *Clin Exp Reprod Med.* 2016; 43(2): 90-96. [CrossRef]
- [14] Ayeleso AO, Oguntibeju OO, Aboua YG, Brooks NL. Effects of red palm oil and rooibos on sperm motility parameters in streptozotocin-induced diabetic rats. *Afr J Tradit Complement Altern Med.* 2014; 11(5): 8-15. [CrossRef]
- [15] Maresch CC, Stute DC, Fleming T, Lin J, Hammes HP, Linn T. Hyperglycemia induces spermatogenic disruption via major pathways of diabetes pathogenesis. *Sci Rep.* 2019; 9(1): 13074. [CrossRef]
- [16] Ghadiri A, Babil FM, Hamidian GR, Oghbaei H, Oskuye ZZ, Ahmadi M, et al. Can troxerutin pretreatment prevent testicular complications in prepubertal diabetic male rats? *Endocr Regul.* 2020; 54(2): 85-95. [CrossRef]

- [17] Shrilatha B, Muralidhara. Occurrence of oxidative impairments, response of antioxidant defences and associated biochemical perturbations in male reproductive milieu in the Streptozotocin-diabetic rat. *Int J Androl.* 2007; 30(6): 508-518. [CrossRef]
- [18] Abbasi Z, Jelodar G, Geramizadeh B. Prevention of diabetic complications by walnut leaf extract via changing aldose reductase activity: An experiment in diabetic rat tissue. *J Diabetes Res.* 2020; 2020: 8982676. [CrossRef]
- [19] Yuluğ E, Türedi S, Karagüzel E, Kutlu Ö, Menteşe A, Alver A. The short term effects of resveratrol on ischemia-reperfusion injury in rat testis. *J Pediatr Surg.* 2014; 49(3): 484-489. [CrossRef]
- [20] Shi GJ, Zheng J, Wu J, Qiao HQ, Chang Q, Niu Y, et al. Beneficial effects of *Lycium barbarum* polysaccharide on spermatogenesis by improving antioxidant activity and inhibiting apoptosis in streptozotocin-induced diabetic male mice. *Food Funct.* 2017;8(3): 1215-1226. [CrossRef]
- [21] Mohamed MZ, Hafez HM, Zenhom NM, Mohammed HH. Cilostazol alleviates streptozotocin-induced testicular injury in rats via PI3K/Akt pathway. *Life Sci.* 2018; 198: 136-142. [CrossRef]
- [22] Szkudelski T. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol Res.* 2001;50(6):537-46.
- [23] Pessoa JC, Etcheverry S, Gambino D. Vanadium compounds in medicine. *Coord Chem Rev.* 2015; 301: 24-48. [CrossRef]
- [24] Kurt O, Ozden TY, Ozsoy N, Tunali S, Can A, Akev N, Yanardag R. Influence of vanadium supplementation on oxidative stress factors in the muscle of STZ-diabetic rats. *Biomaterials.* 2011; 24(5): 943-949. [CrossRef]
- [25] Tian Y, Song W, Xu D, Chen X, Li X, Zhao Y. Autophagy Induced by ROS aggravates testis oxidative damage in diabetes via breaking the feedforward loop linking p62 and Nrf2. *Oxid Med Cell Longev.* 2020; 2020: 7156579. [CrossRef]
- [26] Yuen VG, Orvig C, McNeill JH. Comparison of the glucose-lowering properties of vanadyl sulfate and bis(maltolato)oxovanadium(IV) following acute and chronic administration. *Can J Physiol Pharmacol.* 1995; 73(1): 55-64. [CrossRef]
- [27] Crans DC, Tracey AS. The chemistry of vanadium in aqueous and nonaqueous solution. ACS Symposium Series; American Chemical Society: Washington, DC, 1998. vol 711; pp 2-29. [CrossRef]
- [28] Tunali S, Gezgin-Oktayoglu S, Bolkent S, Coskun E, Bal-Demirci T, Ulkuseven B, Yanardag R. Protective effects of an oxovanadium (IV) complex with N2O2 chelating thiosemicarbazone on small intestine injury of STZ-diabetic rats. *Biol Trace Elem Res.* 2021, 199(4): 1515-1523. [CrossRef]
- [29] Bayrak BB, Tunali S, Bal-Demirci T, Ulkuseven B, Yanardag R. Glycoprotein levels and oxidative lung injury in experimental diabetes: effect of oxovanadium (IV) complex based on thiosemicarbazone. *Toxicol Mech Methods.* 2021, 31(8): 581-588. [CrossRef]
- [30] Wei Y, Zhang C, Zhao P, Yang X, Wang K. A new salicylic acid-derivatized kojic acid vanadyl complex: synthesis, characterization and anti-diabetic therapeutic potential. *J Inorg Biochem.* 2011; 105(8): 1081-1085. [CrossRef]
- [31] Wong CM, Marcocci L, Das D, Wang X, Luo H, Zungu-Edmondson M, et al. Mechanism of protein decarbonylation. *Free Radic Biol Med.* 2013; 65: 1126-1133. [CrossRef]
- [32] Hecker M, Wagner AH. Role of protein carbonylation in diabetes. *JIMD.* 018;41(1):29-38. [CrossRef]
- [33] Bal R, Türk G, Tuzcu M, Yilmaz O, Ozercan I, Kuloglu T, Gür S, Nedzvetsky VS, Tykhomyrov AA, Andrievsky GV, Baydas G, Naziroglu M. Protective effects of nanostructures of hydrated C(60) fullerene on reproductive function in streptozotocin-diabetic male rats. *Toxicology.* 2011; 282(3): 69-81. [CrossRef]
- [34] Lu SC. Regulation of glutathione synthesis. *Mol Aspects Med.* 2009; 30(1-2): 42-59. [CrossRef]
- [35] Vernet P, Aitken RJ, Drevet JR. Antioxidant strategies in the epididymis. *Mol Cell Endocrinol.* 2004; 216: 31-39. [CrossRef]
- [36] Amaral S, Oliveira PJ, Ramalho-Santos J. Diabetes and the impairment of reproductive function: possible role of mitochondria and reactive oxygen species. *Curr. Diabetes Rev.* 2008; 4(1): 46-54. [CrossRef]
- [37] Majd NE, Sadeghi N, Tavalae M, Tabandeh MR, Nasr-Esfahani MH. Evaluation of oxidative stress in testis and sperm of rat following induced varicocele. *Urol J.* 2019; 16(3): 300-306. [CrossRef]

- [38] Khosravi Z, Sedaghat R, Baluchnejadmojarad T, Roghani M. Diosgenin ameliorates testicular damage in streptozotocin-diabetic rats through attenuation of apoptosis, oxidative stress, and inflammation. *Int Immunopharmacol.* 2019; 70: 37-46. [CrossRef]
- [39] Nechay BR, Saunders JP. Inhibition by vanadium of sodium and potassium dependent adenosinetriphosphate derived from animal and human tissues. *J Environ Pathol Toxicol.* 1978; 2(2): 247-262.
- [40] Kustin K. Perspectives on vanadium biochemistry. 1998; Chapter 13; pp. 170-185. [CrossRef]
- [41] Aratani Y. Myeloperoxidase: Its role for host defense, inflammation, and neutrophil function. *Arch Biochem Biophys.* 2018; 640: 47-52. [CrossRef]
- [42] Martínez-Marcos P, Carvajal-Serna M, Lázaro-Gaspar S, Pérez-Pé R, Muiño-Blanco T, Cebrián-Pérez JA, Casao A. Presence of melatonin-catabolizing non-specific enzymes myeloperoxidase and indoleamine 2,3-dioxygenase in the ram reproductive tract. *Reprod Domest Anim.* 2019; 54(12): 1643-1650. [CrossRef]
- [43] Wang SQ, Qin WB, Kang YM, Ma XR, Liu L, Liu JX, Zhang T, Liang Y, Wang F. Intervention effect of ganoderma lucidum spores on the changes of XOD, MPO and SDH in the testis tissue of NIDDM rats. *Zhonghua Nan Ke Xue.* 2008; 14(9): 792-795.
- [44] Liu D, Liu L, Hu Z, Song Z, Wang Y, Chen Z. Evaluation of the oxidative stress-related genes ALOX5, ALOX5AP, GPX1, GPX3 and MPO for contribution to the risk of type 2 diabetes mellitus in the Han Chinese population. *Diab Vasc Dis Res.* 2018; 15(4): 336-339. [CrossRef]
- [45] Yanardag R, Tunali S. Vanadyl sulfate administration protects the streptozotocin-induced oxidative damage to brain tissue in rats. *Mol Cell Biochem.* 2006; 286(1-2): 153-159. [CrossRef]
- [46] Tsounapi P, Honda M, Dimitriadis F, Kawamoto B, Hikita K, Muraoka K, Saito M, Sofikitis N, Takenaka A. Impact of antioxidants on seminal vesicles function and fertilizing potential in diabetic rats. *Asian J Androl.* 2017;19(6):639-646. [CrossRef]
- [47] Tramer F, Rocco F, Micali F, Sandri G, Panfili E. Antioxidant systems in rat epididymal spermatozoa. *Biol Reprod.* 1998; 59(4): 753-758. [CrossRef]
- [48] Schaffer SW, Azuma J, Mozaffari M. Role of antioxidant activity of taurine in diabetes. *Can J Physiol Pharmacol.* 2009;87(2):91-99. [CrossRef]
- [49] Srivastava P, Saxena AK, Kale RK, Baquer NZ. Insulin like effects of lithium and vanadate on the altered antioxidant status of diabetic rats. *Res Commun Chem Pathol Pharmacol.* 1993; 80(3): 283-293.
- [50] Thompson KH, McNeill JH, in: Fischer PWF, L'Abbe MR, Cockell KA, Gibson RS. (Eds.), Trace elements in man and animals 9: Proc. 9th Int. Symp. Trace Elem. Man Animals, NRC Research Press, Ottawa, 1997; pp. 349-350.
- [51] Chandra AK, Ghosh R, Chatterjee A, Sarkar M. Vanadium-induced testicular toxicity and its prevention by oral supplementation of zinc sulphate. *Toxicol Mech Methods.* 2007; 17(4): 175-187. [CrossRef]
- [52] Junod A, Lambert AE, Stauffacher W, Renold AE. Diabetogenic action of streptozotocin: relationship of dose to metabolic response. *J Clin Invest.* 1969; 48(11): 2129-2139. [CrossRef]
- [53] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951; 193(1): 265-275.
- [54] Beutler E. A manual of biochemical methods, vol. 12. London: Academic Press, 1971.
- [55] Ledwozyw A, Michalak J, Stepień A, Kadziolka A. The relationship between plasma triglycerides, cholesterol, total lipids and lipid peroxidation products during human atherosclerosis. *Clin Chim Acta.* 1986; 155(3): 275-283. [CrossRef]
- [56] Levine RL, Garland D, Oliver CN, Amici A, Climent I, Lenz AG, et al. Determination of carbonyl content in oxidatively modified proteins. *Methods Enzymol.* 1990; 186: 464-478. [CrossRef]
- [57] Walter W, Schult C. In: *Methods of Enzymatic Analysis, Vol.2*, Bergmeyer HU, 1974; pp. 856-886.
- [58] Wei H, Frenkel K. In vivo formation of oxidized DNA bases in tumor promoter-treated mouse skin. *Cancer Res.* 1991;51(16):4443-4449.
- [59] Aebi H. Catalase in vitro. *Methods Enzymol.* 1984; 105: 121-126. [CrossRef]

- [60] Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med.* 1967; 70(1): 158-169.
- [61] Wendel A. Glutathione peroxidase. *Methods Enzymol.* 1981; 77: 325-333. [CrossRef]
- [62] Habig WH, Jakoby WB. Assays for differentiation of glutathione S-transferases. *Methods Enzymol.* 1981;77:398-405. [CrossRef]
- [63] Mylroie AA, Collins H, Umbles C, Kyle J. Erythrocyte superoxide dismutase activity and other parameters of copper status in rats ingesting lead acetate. *Toxicol Appl Pharmacol.* 1986; 15; 82(3): 512-520. [CrossRef]