

Relaxing actions of carvacrol on isolated rat duodenum smooth muscles: Evidence for the role of potassium ion channels

Muhammed YUSUF^{1*} , Süleyman AYDIN¹ 

¹ Anadolu University, Institute of Graduate Studies, Faculty of Pharmacy, Department of Pharmacology, Eskişehir, Türkiye

* Corresponding Author. E-mail: ecz.muhammed.yusuf@gmail.com (M.Y.); Tel. 05314508215.

Received: 9 July 2024 / Revised: 27 July 2024 / Accepted: 27 July 2024

ABSTRACT: Mammals have approximately 80 genes that encode potassium (K⁺) channels. They have a wide range of physiological functions, such as modulation of action potential duration and relaxation of smooth muscles. Carvacrol, a compound found in various aromatic plants, has been used traditionally for treating asthma, menstrual spasms, and gastrointestinal disorders. However, its mechanism of action on smooth muscles remains insufficiently understood. This study investigates the role of potassium ion channels in the relaxing effects of carvacrol on isolated rat duodenum smooth muscles. Rats were euthanized and duodenal segments were prepared and placed in an isolated organ bath with Krebs' solution. The segments were equilibrated for 1 h and then treated with chemicals to obtain concentration-response curves. The data were evaluated using one-way Analysis of Variance (ANOVA) and Tukey's Honest Significant Difference (Tukey's HSD) test for multiple comparisons. Our findings demonstrate that carvacrol induces a dose-dependent relaxation of these muscles. The relaxation effects were significantly reduced in the presence of specific potassium channel inhibitors, including paxilline, UCL1684, linopirdine, barium chloride, and 4-aminopyridine. They were completely blocked by the combination of barium chloride and tetraethylammonium. However, glibenclamide, ruthenium red, and nitroarginine did not alter the relaxing effects of carvacrol. In conclusion: carvacrol relaxes duodenum smooth muscles by opening barium chloride-sensitive Kir2.1 and Kir3.1, as well as tetraethylammonium-sensitive KCa1.1, KCa2.1, KCa2.2, KCa2.3, Kv1.2, Kv1.4, Kv1.5, Kv2.1, Kv2.2, Kv4.1, Kv4.2, Kv4.3, Kv7.1, Kv7.2, Kv7.3, Kv7.4, and Kv7.5 potassium channels.

KEYWORDS: Carvacrol; Ion channel; Transient receptor potential; Potassium; Duodenum; Smooth muscle.

1. INTRODUCTION

Changes in the contractility of smooth muscles play an important role in the physiopathology of various diseases such as irritable bowel syndrome, achalasia, and hyperactive bladder [1-3]. Mammals have approximately 80 genes that encode potassium (K⁺) channels [4]. The four primary types of human K⁺ channels are inward-rectifying (Kir), calcium-activated (KCa), two-pore (K2P), and voltage-gated (Kv) channels [5]. They are expressed in numerous tissues and have a wide range of physiological functions, such as modulation of action potential duration and relaxation of smooth muscle cells [6]. Among the potassium channels that are distributed and play a role in the relaxation of smooth muscles are KCa1.1, KCa2.1, KCa2.2, KCa2.3, Kv2.1, Kv2.2, Kv4.1, Kv4.2, Kv4.3, Kv7.1, Kv7.2, Kv7.3, Kv7.4, Kv7.5, Kv1.2, Kv1.5, Kv1.4, Kir2.1, Kir3.1, Kir6.1, and Kir6.2 [4,7].

Carvacrol (2-methyl-5-(1-methyl ethyl) phenol) is a synthetic chemical compound that can also be naturally found in various aromatic plants [8]. Since ancient times, carvacrol-containing plants have been used ethnomedically for various purposes, especially against asthma, menstrual spasms, and gastrointestinal disorders [8]. In studies on smooth muscles, it has been reported that carvacrol relaxes the smooth muscles of the trachea independently of beta-adrenergic, histaminergic, and muscarinic receptor interactions [9]. However, subsequent research has demonstrated that carvacrol exhibits antimuscarinic and antihistaminic effects [10,11] and acts as an adrenergic agonist on the trachea [12]. On the other hand, it has been reported that carvacrol is ineffective on the aorta [13]. However, other studies have shown that it relaxes the aorta and mesenteric artery, likely by reducing calcium release from the sarcoplasmic reticulum and calcium entry from the cell membrane and transient receptor potential (TRP) ion channels [14,15].

How to cite this article: Yusuf M, Aydin S. Relaxing actions of carvacrol on isolated rat duodenum smooth muscles: evidence for the role of potassium ion channels. *J Res Pharm.* 2024; 28(5): 1501-1507.

Considering the studies conducted to date with carvedilol on smooth muscles and the diversity of potassium ion channels involved in smooth muscle relaxation, the mechanism of action of carvedilol remains insufficiently clarified. Therefore, this study was conducted to elucidate the role of potassium channels in the mechanisms mediating carvedilol's effect on the smooth muscles of the rat duodenum.

2. RESULTS

In our study, carvedilol showed relaxant effects on the smooth muscles of the duodenum in a dose-dependent manner (Figures 1, 2, 3). The relaxing effect of carvedilol decreased in the presence of paxilline, UCL1684, linopirdine (Figure 1) (Table 1), barium chloride (BaCl₂) (Figure 2) (Table 2), and 4-aminopyridine (4-AP). (Figure 3) (Table 3). The relaxant effect of carvedilol was completely blocked in the presence of BaCl₂ + tetraethylammonium (TEA) (Figure 2) (Table 2). In contrast, in the presence of glibenclamide (Figure 2) (Table 2), ruthenium red, and nitroarginine (L-NOARG) (Figure 3) (Table 3), there was no change in the relaxant effect of carvedilol.




Figure 1. Effect of paxilline, UCL1684, and linopirdine on the relaxant effect of carvedilol on isolated rat duodenal smooth muscles. Maximum relaxation by carvedilol was reduced in the presence of paxilline, UCL1684, and linopirdine. (n=7). $p < 0.05$ (*), 0.01 (**), 0.001 (***)

Table 1. Comparison of the IC₅₀ values for the effects of paxillin, UCL1684, and linopirdine on the relaxing effect of carvedilol on rat duodenum smooth muscles.

Groups	IC ₅₀ (estimate)	IC ₅₀ (SEM)	p
Carvedilol / DMSO	1.6284e+00	1.0054e+01	0.95029 ***
Carvedilol / Paxilline+Carvedilol	2.9720e+03	2.5179e+03	0.24080 ***
Carvedilol / UCL1684+Carvedilol	1.7390e+02	1.2160e+03	0.88722 *
Carvedilol / Linopirdine+Carvedilol	1.8041e-01	4.7643e-02	< 2e-16 *
DMSO / Paxilline+Carvedilol	1.8251e+03	1.1360e+04	0.87275
DMSO / UCL1684+Carvedilol	1.0680e+02	9.9565e+02	0.91559
DMSO / Linopirdine+Carvedilol	1.1079e-01	6.8377e-01	0.19644
Paxilline+Carvedilol / UCL1684+Carvedilol	5.8513e-02	4.1173e-01	-0.02433 *
Paxilline+Carvedilol / Linopirdine+Carvedilol	6.0702e-05	5.0345e-05	< 2e-16 ***
UCL1684+Carvedilol / Linopirdine+Carvedilol	1.0374e-03	7.2518e-03	< 2e-16 ***

SEM: standard error of mean; $p < 0.05$ (*), 0.01 (**), 0.001 (***)




Figure 2. Effect of BaCl₂, TEA + BaCl₂, and glibenclamide on the relaxant effect of carvacrol on isolated rat duodenal smooth muscles. Maximum relaxation by carvacrol was reduced in the presence of BaCl₂ and was completely blocked in the presence of BaCl₂ + TEA, in the presence of glibenclamide there was no change in the relaxant effect of carvacrol (n=7). *p*<0.05 (*), 0.01 (**), 0.001 (***).

Table 2. Comparison of the IC₅₀ values for the effects of BaCl₂, TEA +BaCl₂, and glibenclamide on the relaxing effect of carvacrol on rat duodenum smooth muscles

Groups	IC ₅₀ (estimate)	IC ₅₀ (SEM)	p
Carvacrol / DMSO	1.6284e+00	1.0054e+01	0.95029 ***
Carvacrol / BaCl ₂ +Carvacrol	2.9720e+03	2.5179e+03	0.24080 *
Carvacrol / TEA+BaCl ₂ +Carvacrol	1.7390e+02	1.2160e+03	0.88722 ***
Carvacrol / Glibenclamide+Carvacrol	1.8041e-01	4.7643e-02	< 2e-16
DMSO / BaCl ₂ +Carvacrol	1.8251e+03	1.1360e+04	0.87275
DMSO / TEA+BaCl ₂ +Carvacrol	1.0680e+02	9.9565e+02	0.91559
DMSO / Glibenclamide+Carvacrol	1.1079e-01	6.8377e-01	0.19644 * *
BaCl ₂ +Carvacrol / TEA+BaCl ₂ +Carvacrol	5.8513e-02	4.1173e-01	0.02433 *
BaCl ₂ +Carvacrol / Glibenclamide+Carvacrol	6.0702e-05	5.0345e-05	< 2e-16 *
TEA+BaCl ₂ +Carvacrol / Glibenclamide+Carvacrol	1.0374e-03	7.2518e-03	< 2e-16 ***

SEM: standard error of mean; *p*<0.05 (*), 0.01 (**), 0.001 (***).




Figure 3. Effect of 4-AP, ruthenium red, and L-NOARG on the relaxant effect of carvacrol on isolated rat duodenal smooth muscles. Maximum relaxation by carvacrol was reduced in the presence of 4-AP, in the presence of ruthenium red, and L-NOARG there was no change in the relaxant effect of carvacrol ($n=7$). $p<0.05$ (*), 0.01 (**), 0.001 (***)

Table 3. Comparison of the IC_{50} values for the effects of 4-AP, ruthenium red, and L-NOARG on the relaxing effect of carvacrol on rat duodenum smooth muscles.

Groups	IC_{50} (estimate)	IC_{50} (SEM)	p
Carvacrol/ DMSO	1.093057	9.381999	0.9921 ***
Carvacrol / 4-AP+Carvacrol	0.323375	0.527365	<2e-16 ***
Carvacrol / Ruthenium Red+Carvacrol	2.254434	1.205454	0.3006
Carvacrol / L-NOARG+Carvacrol	0.089918	0.071585	0.2024
DMSO / 4-AP+Carvacrol	0.295845	2.576608	0.7852
DMSO / Ruthenium Red+Carvacrol	2.062504	17.679743	0.9522 ***
DMSO / L-NOARG+Carvacrol	0.082263	0.706822	0.1971 ***
4-AP+Carvacrol / Ruthenium Red+Carvacrol	6.971580	10.947604	0.5866 **
4-AP+Carvacrol / L-NOARG+Carvacrol	0.278061	0.466429	0.1248 **
Ruthenium Red+Carvacrol /L-NOARG+Carvacrol	0.039885	-36.2845	0.6896

SEM: standard error of mean; $p<0.05$ (*), 0.01 (**), 0.001 (***)

3. DISCUSSION

It has been reported that carvacrol affected potassium channels in the substantia gelatinosa of the spinal medulla, which were not sensitive to barium and tetraethylammonium [16]. However, in our studies, the relaxing effect of carvacrol was diminished in the presence of various channel inhibitors; BaCl₂ (Kir2.1 and Kir3.1) (Figure 2), 4-AP (Kv1.2, Kv1.4, Kv1.5, Kv2.1, Kv2.2, Kv4.1, Kv4.2, and Kv4.3) (Figure 3), linopirdine (Kv7.1, Kv7.2, Kv7.3, Kv7.4, and Kv7.5) (Figure 1), paxillin (KCa1.1) (Figure 1), and UCL1684 (KCa2.1, KCa2.2, and KCa2.3) (Figure 1); and was completely blocked in the presence of BaCl₂ (Kir2.1 and Kir3.1) + TEA (Kv1.2, Kv1.4, Kv1.5, Kv2.1, Kv2.2, Kv4.1, Kv4.2, Kv4.3, Kv7.1, Kv7.2, Kv7.3, Kv7.4, Kv7.5, KCa1.1, KCa2.1, KCa2.2, and KCa2.3) (Figure 2). Our findings indicate that carvacrol interacts with barium chloride -sensitive; Kir2.1, Kir3.1, and tetraethylammonium-sensitive; KCa1.1, KCa2.1, KCa2.2, KCa2.3, Kv1.2, Kv1.4, Kv1.5, Kv2.1, Kv2.2, Kv4.1, Kv4.2, Kv4.3, Kv7.1, Kv7.2, Kv7.3, Kv7.4, Kv7.5 potassium channels. Compared to previously reported findings, differences between neurons and smooth muscles may lead to different and even contradictory experimental results. These results suggest that carvacrol does not exhibit the same effects on every tissue and cell, and its effects vary on different tissues and cell types.

Previous studies reported that carvacrol protects against oxidative damage via Kir6.2 channels [17]. However, in our studies, it was observed that the relaxing effect of carvacrol did not change in the presence of glibenclamide, an inhibitor of Kir6.1 and Kir6.2 channels (Figure 2). Contrary to previously reported findings, our findings indicate that Kir6.1 and Kir6.2 channels do not play any role in the relaxing effect of carvacrol observed on the gastrointestinal system's smooth muscles, and the effect of carvacrol in the oxidative damage mechanism may differ from its effect on smooth muscles.

The non-adrenergic non-cholinergic (NANC) system, which uses nitric oxide (NO) as a transmitter, is a part of the autonomic nervous system with a key role in gastrointestinal relaxations [18]. Some studies reported that the gastroprotective effect of carvacrol is mediated by the activation of nitric oxide synthase [19]. In our study, the lack of blocking effect through the nitric oxide synthase inhibitor L-NOARG (Figure 3), indicates no significant role for nitric oxide in the relaxant effect of carvacrol on the duodenum smooth muscles.

Transient Receptor Potential ion channels include at least 28 subtypes in mammals [20]; and they are vital in regulating gastrointestinal (GI) smooth muscle relaxation and contraction. The key TRP channels involved include TRPC, TRPV, and TRPM [21, 22]. Some of them play an important role in transferring calcium and sodium ions into the cell [23, 24]. Different studies have reported the role of TRP channels in the mechanism of action of carvacrol; and while some of them are activated (TRPV3 and TRPA1) [25, 26], others are inhibited by carvacrol (TRPC1 and TRPM7) [27, 28]. The lack of effect of ruthenium red (Figure 3), indicates no significant role for ruthenium red-sensitive TRP ion channels in the relaxant effect of carvacrol on the duodenum smooth muscles.

Potassium channels are not just an ion channel family belonging to humans and mammals. They are an important ion channel family that appeared very early in the evolutionary development of life on Earth and are found in almost all living organisms, from protozoa to single-celled organisms [29]. Our findings suggest that KCa1.1, KCa2.1, KCa2.2, KCa2.3, Kv1.2, Kv1.4, Kv1.5, Kv2.1, Kv2.2, Kv4.1, Kv4.2, Kv4.3, Kv7.1, Kv7.2, Kv7.3, Kv7.4, Kv7.5, Kir2.1, and Kir3.1 channels play a major role in the relaxing effect of carvacrol on isolated duodenum smooth muscles. Furthermore, a review of the UniProt protein database as of December 2023 (version 2023_05) revealed 90 potassium channel proteins and 92 isoforms in humans (<http://ftp.ebi.ac.uk/>). Currently, there are no specific agonists or antagonists available for each of these ion channels; and due to the large number and variety of these channels and the lack of sufficiently specific agonists and/or antagonists for each of these channels, further research is needed on the effects of carvacrol on potassium channels.

4. CONCLUSION

Carvacrol relaxes duodenum smooth muscles by opening barium chloride and tetraethylammonium-sensitive potassium channels. Our research highlights its effects on a broad range of potassium channels and suggests it is a noteworthy subject for further study.

5. MATERIALS AND METHODS

5.1. Chemicals

Carvacrol, paxilline, ruthenium red, L-NOARG, UCL1684, linopirdine, TEA, 4-aminopyridine, glibenclamide, BaCl₂, dimethyl sulfoxide (DMSO), ketamine hydrochloride, KCl, NaCl, NaHCO₃, d-glucose, KH₂PO₄, MgSO₄·7H₂O, and CaCl₂·2H₂O were purchased from Sigma-Aldrich (Germany).

5.2. Animals

Ethical approval for the study was obtained from the ethics committee of Anadolu University on 16.06.2021 (Decision no: 2021-36). The study utilized male Sprague-Dawley rats, each weighing 220-250 g, obtained from the Experimental Animals Research Center at Anadolu University. These rats were maintained in a controlled environment with a 12-h light/dark cycle, 55%-60% humidity, and a constant temperature of 22 °C. They had ad libitum access to food and water.

5.3. Isolated duodenum experiments

After anesthetizing the rats with ketamine hydrochloride, they were euthanized by cervical dislocation. The duodenal organs were excised and placed in Krebs' solution with the following composition (in mM): NaCl, 118.4; KCl, 4.7; MgSO₄·7H₂O, 1.2; CaCl₂·2H₂O, 2.5; NaHCO₃, 25; KH₂PO₄, 1.2; and d-glucose 11. The organs were cleaned of fat and connective tissue, and 1-1.5 cm segments were prepared. These segments were placed in isolated organ baths (Ugo Basile, Varese, Italy) with 20 ml of Krebs' solution (pH

7.4), continuously aerated with 5% CO₂ and 95% O₂ at 37°C. The segments were connected to isotonic transducers which were connected to a two-channel pen recorder. The tissues were equilibrated for 1 h, during this period the solution was replaced every 15 minutes while maintaining a resting tension of 1 g. Then, chemicals were applied to the tissues, and cumulative concentration-response curves were obtained. In the experiments, the following substances were used: carvacrol (10⁻⁷, 10⁻⁶, 10⁻⁵, 10⁻⁴ M), dissolved in water and DMSO (1:1) [13]; and paxilline (10⁻⁶ M) [30], UCL1684 (10⁻⁷ M) [31], linopirdine (10⁻⁶ M) [32], tetraethylammonium (10⁻² M) [33], 4-aminopyridine (10⁻³ M) [33], glibenclamide (10⁻⁶ M) [34], BaCl₂ (10⁻³ M) [35], ruthenium red (10⁻⁵ M) [36], and L-NOARG (10⁻⁴ M) [37], all dissolved in water.

5.4. Statistical analysis of data

The R programming language and related packages were used for statistical evaluation of experimental data, nonlinear regression, and graphics drawing (R core team, 2019). The data were evaluated using one-way analysis of variance (ANOVA) and Tukey's HSD test for multiple comparisons. The IC₅₀ was determined by non-linear regression analysis of log-dose vs. response curves. A p value <0.05 was considered as statistically significant. Data is expressed as the estimate ± standard error of mean [SEM].

Acknowledgments: This article is a study produced from the doctoral thesis of Muhammed YUSUF (Anadolu University, Institute of Graduate Studies).

Author contributions: Concept - M.Y., S.A.; Design - M.Y., S.A.; Supervision - S.A.; Resources - M.Y.; Materials - M.Y.; Data Collection and/or Processing - M.Y.; Analysis and/or Interpretation - M.Y., S.A.; Literature Search - M.Y., S.A.; Writing - M.Y.; Critical Reviews - M.Y., S.A.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Tack, J, Talley, NJ, Camilleri, M, Holtmann, G, Hu, P, Malagelada, JR. Functional gastroduodenal disorders. *Gastroenterol.* 2006; 130 (5): 1466-1479. <https://doi.org/10.1053/j.gastro.2005.11.059>
- [2] Lacy BE, Weiser K. Gastrointestinal motility disorders: an update. *Dig Dis.* 2006; 24 (3-4): 228-242. <https://doi.org/10.1159/000092876>
- [3] Andersson KE, Arner A. Urinary bladder contraction and relaxation: Physiology and pathophysiology. *Physiol Rev.* 2004; 84 (3): 935-986. <https://doi.org/10.1152/physrev.00038.2003>
- [4] Curro D. The modulation of potassium channels in the smooth muscle as a therapeutic strategy for disorders of the gastrointestinal tract. *Adv Protein Chem Struct Biol.* 2016; 104: 263-305. <https://doi.org/10.1016/bs.apcsb.2015.12.002>
- [5] Maljevic S, Lerche H. Potassium channels: A review of broadening therapeutic possibilities for neurological diseases. *J Neurol.* 2013; 260 (9): 2201-2211. <https://doi.org/10.1007/s00415-012-6727-8>
- [6] Vyas VK, Parikh P, Ramani J, Ghate M. Medicinal chemistry of potassium channel modulators: An update of recent progress (2011-2017). *Curr Med Chem.* 2019; 26 (12): 2062-2084. <https://doi.org/10.2174/0929867325666180430152023>
- [7] Beyder A, Farrugia G. Targeting ion channels for the treatment of gastrointestinal motility disorders. *Therap Adv Gastroenterol.* 2012; 5 (1): 5-21. <https://doi.org/10.1177/1756283x11415892>
- [8] Baser KHC. Biological and pharmacological activities of carvacrol and carvacrol bearing essential oils. *Curr Pharm Des.* 2008; 14(29): 3106-3120. <https://doi.org/10.2174/138161208786404227>
- [9] Boskabady MH, Jandaghi P. Relaxant effects of carvacrol on guinea pig tracheal chains and its possible mechanisms. *Pharmazie.* 2008; 58 (9): 661-663.
- [10] Boskabady MH, Jafari Z, Pouraboli I. The effect of carvacrol on muscarinic receptors of guinea-pig tracheal chains. *Phytother Res.* 2011; 25 (4): 530-535. <https://doi.org/10.1002/ptr.3290>
- [11] Boskabady MH, Tabanfar H, Gholamnezhad Z, Sadeghnia HR. Inhibitory effect of *Zataria multiflora* Boiss and carvacrol on histamine (H₁) receptors of guinea-pig tracheal chains. *Fundam Clin Pharmacol.* 2012; 26 (5): 609-620. <https://doi.org/10.1111/j.1472-8206.2011.00971.x>
- [12] Boskabady MH, Kaveh M, Eftekhari N, Nemati A. *Zataria multiflora* Boiss and carvacrol affect β₂-adrenoceptors of guinea pig trachea. *Evid Based Complement Alternat Med.* 2010; 857124. <https://doi.org/10.1155/2011/857124>
- [13] Aydin Y, Kutlay O, Ari S, Duman S, Uzuner K, Aydin S. Hypotensive effects of carvacrol on the blood pressure of normotensive rats. *Planta Med.* 2007; 73 (13): 1365-1371. <https://doi.org/10.1055/s-2007-990236>
- [14] Dantas BP, Alves QL, Assis KS, Ribeiro TP, Almeida MM, Vasconcelos AP, Araujo DAM, Braga VA, Medeiros IA, Alencar JL, Silva DF. Participation of the TRP channel in the cardiovascular effects induced by carvacrol in normotensive rat. *Vascul Pharmacol.* 2015; 67-69: 48-58. <https://doi.org/10.1016/j.vph.2015.02.016>
- [15] Peixoto-Neves D, Silva-Alves KS, Gomes MD, Lima FC, Lahlou S, Magalhaes PJ, Ceccatto VM, Coelho-de-Souza AN, Leal-Cardoso JH. Vasorelaxant effects of the monoterpene phenol isomers, carvacrol, and thymol, on rat isolated aorta. *Fundam Clin Pharmacol.* 2010; 24 (3): 341-350. <https://doi.org/10.1111/j.1472-8206.2009.00768.x>

- [16] Luo QT, Fujita T, Jiang CY, Kumamoto E. Carvedilol presynaptically enhances spontaneous excitatory transmission and produces outward current in adult rat spinal substantia gelatinosa neurons. *Brain Res.* 2014; 1592: 44-54. <https://doi.org/10.1016/j.brainres.2014.10.021>
- [17] Balci CN, Firat T, Acar N, Kukner A. Carvedilol treatment opens Kir6.2 ATP-dependent potassium channels and prevents apoptosis on rat testis following ischemia-reperfusion injury model. *Rom J Morphol Embryol.* 2021; 62 (1): 179-190. <https://doi.org/10.47162/rjme.62.1.17>
- [18] Komori S, Unno T, Ohashi H. Pharmacological properties of non-adrenergic, non-cholinergic inhibitory transmission in chicken gizzard. *Br J Pharmacol.* 1997; 121 (3): 572 - 578. <https://doi.org/10.1038%2Fsj.bjp.0701132>
- [19] Oliveira IS, Silva FV, Viana AFSC. Gastroprotective activity of carvedilol on experimentally induced gastric lesions in rodents. *Naunyn Schmiedebergs Arch Pharmacol.* 2012; 385 (9): 899-908. <https://doi.org/10.1007/s00210-012-0771-x>
- [20] Nilius B, Owsianik G. The transient receptor potential family of ion channels. *Gen Biol.* 2011; 12 (3): 218. <https://doi.org/10.1186/gb-2011-12-3-218>
- [21] Earley S, Heppner TJ, Nelson MT, Brayden JE. TRPV4 forms a novel Ca²⁺ signaling complex with ryanodine receptors and BKCa channels. *Circ Res.* 2005; 97 (12): 1270-1279. <https://doi.org/10.1161/01.res.0000194321.60300.d6>
- [22] Johnson CD, Melanaphy D, Purse A, Stokes NR, Dickenson AH. TRPM8 channel as a therapeutic target. *Am J Physiol Heart Circ Physiol.* 2009; 296 (6): 522-532. <https://doi.org/10.1152/ajpheart.01112.2008>
- [23] Beech DJ. TRPC1: Store-operated channel and more. *Pflugers Arch.* 2005; 451 (1): 53-60. <https://doi.org/10.1007/s00424-005-1441-3>
- [24] Veldhuis NA, Poole D, Grace M, McIntyre P, Bunnett NW. The G protein-coupled receptor-transient receptor potential channel axis: Molecular insights for targeting disorders of sensation and inflammation. *Pharmacol Rev.* 2015; 67 (1): 36-73. <https://doi.org/10.1124/pr.114.009555>
- [25] Xu H, Delling M, Jun JC, Clapham DE. Oregano, thyme and clove-derived flavors and skin sensitizers activate specific TRP channels. *Nat Neurosci.* 2006; 9(5): 628-635. <https://doi.org/10.1038/nm1692>
- [26] Saito S, Nakatsuka K, Takahashi K, Fukuta N, Imagawa T, Ohta T, Tominaga M. Analysis of transient receptor potential ankyrin 1 (TRPA1) in frogs and lizards illuminates both nociceptive heat and chemical sensitivities and coexpression with TRP vanilloid 1 (TRPV1) in ancestral vertebrates. *J Biol Chem.* 2012; 287 (37): 30743-30754. <https://doi.org/10.1074%2Fjbc.M112.362194>
- [27] Peters M, Trembovler V, Alexandrovich A, Parnas M, Birnbaumer L, Minke B, Shohami E. Carvedilol together with TRPC1 elimination improve functional recovery after traumatic brain injury in mice. *J Neurotrauma.* 2012; 29 (18): 2831-2834. <https://doi.org/10.1089%2Fneu.2012.2575>
- [28] Khalil A, Kovac G, Walker MC. Carvedilol after status epilepticus (SE) prevents recurrent SE, early seizures, cell death, and cognitive decline. *Epilepsia.* 2017; 58: 263-273.
- [29] Jegla T, Salkoff L. Molecular evolution of K⁺ channels in primitive eukaryotes. *Soc Gen Physiol Ser.* 1994; 49: 213-222.
- [30] Xin W, Li N, Cheng, Petkov GV. BK Channel-mediated relaxation of urinary bladder smooth muscle: A novel paradigm for phosphodiesterase type 4 regulation of bladder function. *J Pharmacol Exp Ther.* 2014; 349 (1): 56-65. <https://doi.org/10.1124%2Fjpet.113.210708>
- [31] Karpinska O, Baranowska-Kuczko M, Malinowska B, Kloza M, Kusaczuk M, Gegotek A, Golec P, Kasacka I, Kozłowska H. Mechanisms of l- α -lysophosphatidylinositol-induced relaxation in human pulmonary arteries. *Life Sci.* 2018; 192: 38-45. <https://doi.org/10.1016/j.lfs.2017.11.020>
- [32] Sakakibara K, Feng G, Li J, Akahori T, Yasuda Y, Nakamura E, Hatakeyama N, Fujiwara Y, Kinoshita H. Kynurenine causes vasodilation and hypotension induced by activation of KCNQ-encoded voltage-dependent K⁺ channels. *J Pharmacol Sci.* 2015; 129 (1): 31-37. <https://doi.org/10.1016/j.jphs.2015.07.042>
- [33] Dong D, Wang Q, Chen W, Fan J, Mu J, Ke J, Yang B. Contrasting effects of tetraethylammonium and 4-aminopyridine on the gastrointestinal function of mice. *Eur J Pharmacol.* 2005; 509 (2-3):179-85. <https://doi.org/10.1016/j.ejphar.2005.01.013>
- [34] Zhang H, Bolton TB. Activation by intracellular GDP, metabolic inhibition and pinacidil of a glibenclamide-sensitive K-channel in smooth muscle cells of rat mesenteric artery. *Br J Pharmacol.* 1995; 114 (3): 662-672. <https://doi.org/10.1111/j.1476-5381.1995.tb17190.x>
- [35] Huang Y. BaCl₂- and 4-aminopyridine-evoked phasic contractions in the rat vas deferens. *Br J Pharmacol.* 1995; 115 (5): 845-851. <https://doi.org/10.1111%2Fj.1476-5381.1995.tb15010.x>
- [36] Nozawa K, Kawabata-Shoda E, Doihara H, Kojima R, Okada H, Mochizuki S, Sano Y, Inamura K, Matsushime H, Koizumi T, Yokoyama T, Ito H. TRPA1 regulates gastrointestinal motility through serotonin release from enterochromaffin cells. *Proc Natl Acad Sci USA.* 2009; 106: 3408-3413. <https://doi.org/10.1073/pnas.0805323106>
- [37] Yuan SY, Brookes SJ, Costa M. Distension-evoked ascending and descending reflexes in the isolated guinea-pig stomach. *J Auton Nerv Syst.* 1997; 62(1-2): 94-102. [https://doi.org/10.1016/s0165-1838\(96\)00115-4](https://doi.org/10.1016/s0165-1838(96)00115-4)

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dspace.marmara.edu.tr>.